

SL₂ QUANTUM TRACE IN QUANTUM TEICHMÜLLER THEORY VIA WRITHE

HYUN KYU KIM, THANG T. Q. LÊ, AND MIRI SON

ABSTRACT. Quantization of Teichmüller space of a punctured Riemann surface S is an approach to three dimensional quantum gravity, and is a prototypical example of quantization of cluster varieties. Any simple loop γ in S gives rise to a natural trace-of-monodromy function $\mathbb{I}(\gamma)$ on the Teichmüller space. For any ideal triangulation Δ of S , this function $\mathbb{I}(\gamma)$ is a Laurent polynomial in the square-roots of the exponentiated shear coordinates for the arc of Δ . An important problem was to construct a quantization of this function $\mathbb{I}(\gamma)$, namely to replace it by a non-commutative Laurent polynomial in the quantum variables. This problem, which is closely related to the framed protected spin characters in physics, has been solved by Allegretti and Kim using Bonahon-Wong’s SL₂ quantum trace for skein algebras, and by Gabella using Gaiotto-Moore-Neitzke’s Seiberg-Witten curves, spectral networks, and writhe of links. We show that these two solutions to the quantization problem coincide. We enhance Gabella’s solution and show that it is a twist of Bonahon-Wong’s quantum trace.

CONTENTS

1. Introduction	1
1.1. Quantization of Teichmüller space	1
1.2. Allegretti-Kim quantization using Bonahon-Wong quantum trace	2
1.3. Another quantization using Gabella quantum holonomy	3
1.4. More general result, and proof	4
2. Quantum Teichmüller theory	5
2.1. Ideal triangulation of a non-compact surface	5
2.2. Teichmüller space and shear coordinates	8
2.3. Quantum Teichmüller space	9
2.4. Deformation quantization of Teichmüller space and the quantum ordering problem	10
3. Bonahon-Wong’s quantum trace	11
3.1. Tangles and skein algebra	11
3.2. Square-root quantum Teichmüller space	14
3.3. Quantum trace map	16
3.4. Biangle quantum trace	19
3.5. State-sum formula	21
4. Gabella’s quantum holonomy	23
4.1. Branched double cover surface	23
4.2. Biangle quantum holonomy via operator invariant	26
4.3. Quantum holonomy	29
5. Equality of the two constructions	32
5.1. Statement of the main theorem	32
5.2. Bonahon-Wong triangle factors	34
5.3. Gabella triangle factors	34
5.4. Bonahon-Wong biangle factors via $\mathcal{U}_q(\mathfrak{sl}_2)$ Reshetikhin-Turaev operator invariant	37
5.5. Gabella biangle factors as twisting of Bonahon-Wong biangle factors	40
5.6. Finishing the proof of main theorem	43
References	43

1. INTRODUCTION

1.1. Quantization of Teichmüller space. Quantization of the Teichmüller space of a Riemann surface appeared in late 1990’s as an approach to (2+1)-quantum gravity in mathematical physics. Certain versions of quantum Teichmüller spaces were first established by Kashaev [K98] and independently by Chekhov and Fock [CF99] [F97]. The Chekhov-Fock formulation was later generalized by Fock and

Goncharov to the theory of quantum cluster varieties [FG09]. Here we review the Chekhov-Fock(-Goncharov) quantum Teichmüller theory.

Let \mathfrak{S} be a *punctured surface*, which is the result of removing a finite number of points, called *punctures*, from a closed oriented surface of finite genus. Among several versions of Teichmüller spaces of \mathfrak{S} , we will consider the *enhanced Teichmüller space* $\mathcal{X}^+(\mathfrak{S})$, also known as the *holed Teichmüller space* [F97] [FG07]. By definition, $\mathcal{X}^+(\mathfrak{S})$ is the set of all complete hyperbolic metrics h on \mathfrak{S} up to isotopy, enhanced with a choice of an orientation of a small loop surrounding each puncture whose monodromy with respect to h is hyperbolic. The space $\mathcal{X}^+(\mathfrak{S})$ is a contractible smooth manifold equipped with the *Weil-Petersson* Poisson structure [F97] [FG07]. A nice coordinate system requires the choice of an *ideal triangulation* Δ of \mathfrak{S} , which is a collection of isotopy classes of mutually non-intersecting simple paths running between punctures, called (*ideal*) *arcs*, dividing \mathfrak{S} into (*ideal*) *triangles*. Per each arc e of Δ , Thurston's *shear coordinate* function $x_e = x_{e;\Delta}$ is associated [T80], and they together comprise a global coordinate system of $\mathcal{X}^+(\mathfrak{S})$. Their exponentials $X_e := \exp(x_e)$ are particularly convenient. The Poisson bracket is given by $\{X_e, X_f\} = \varepsilon_{ef} X_e X_f$, $\forall e, f \in \Delta$, where $\varepsilon_{ef} = a_{ef} - a_{fe} \in \mathbb{Z}$, with a_{ef} being the number of corners of ideal triangles of Δ formed by e and f appearing clockwise this order. Per change of ideal triangulations, the coordinate change formulas for the exponentiated shear coordinates are rational, providing examples of so-called cluster \mathcal{X} -mutations [FG06].

For a chosen triangulation Δ , a quantum deformed version of the classical Poisson algebra generated by the exponentiated shear coordinates X_e and their inverses had been constructed as the \mathbb{C} -algebra generated by the symbols \widehat{X}_e and their inverses, with e running through all arcs of Δ , mod out by the relations $\widehat{X}_e \widehat{X}_f = q^{2\varepsilon_{ef}} \widehat{X}_f \widehat{X}_e$, $\forall e, f \in \Delta$, where $q \in \mathbb{C}^*$ is the quantum parameter; this algebra is called the *Chekhov-Fock algebra* \mathcal{T}_Δ^q [L09]. A key aspect of quantum Teichmüller theory is consistency under change of triangulations. It has been shown that, for each pair of ideal triangulations Δ and Δ' , there exists a quantum coordinate change isomorphism $\Phi_{\Delta, \Delta'}^q : \text{Frac}(\mathcal{T}_{\Delta'}^q) \rightarrow \text{Frac}(\mathcal{T}_\Delta^q)$ between the skew-fields of fractions of the Chekhov-Fock algebras, that recovers the classical formula as $q = 1$ and satisfies the consistency equations $\Phi_{\Delta, \Delta''}^q = \Phi_{\Delta, \Delta'}^q \circ \Phi_{\Delta', \Delta''}^q$ [CF99] [L09]. The algebras $\text{Frac}(\mathcal{T}_\Delta^q)$ for different Δ 's can then be identified by these isomorphisms in a consistent manner, forming a *quantum Teichmüller space* [L09].

The algebraic aspects of [CF99] [F97] [K98] [FG09] were mostly on establishing this consistent quantum Teichmüller space as just described, but not so much on finding a *deformation quantization map*, i.e. a map sending a classical function to a quantum counterpart. One must first determine which functions to quantize. Fock and Goncharov [FG06] suggested to consider the functions that can be written as Laurent polynomials in the exponentiated shear coordinates $(X_e)_{e \in \Delta}$ for *every* ideal triangulation Δ , i.e. the regular functions on $\mathcal{X}^+(\mathfrak{S})$ viewed as $(\mathbb{R}_{>0}$ -points of) a Fock-Goncharov cluster \mathcal{X} -variety. This important class of functions called the *universally Laurent functions* form the classical Poisson algebra to be quantized, which we denote by $\mathbf{L}(\mathcal{X}^+(\mathfrak{S}))$. By a deformation quantization we mean a map

$$(1.1) \quad \mathbb{I}_\Delta^q : \mathbf{L}(\mathcal{X}^+(\mathfrak{S})) \rightarrow \mathcal{T}_\Delta^q$$

satisfying favorable properties, e.g. when $q = 1$ this should be the tautological classical map, and this map (for general q) must be consistent under the quantum coordinate change $\mathbb{I}_{\Delta'}^q = \mathbb{I}_\Delta^q \circ \Phi_{\Delta, \Delta'}^q$. Existence of such a map was conjectured in [FG06] and a solution was first given by Allegretti and the first author in [AK17]. Later, another solution is given by Gabella [G17]. The goal of the present paper is to compare these two solutions.

1.2. Allegretti-Kim quantization using Bonahon-Wong quantum trace. A basic example of a universally Laurent function on the enhanced Teichmüller space $\mathcal{X}^+(\mathfrak{S})$ is the trace-of-monodromy function along a loop. Let γ be any non-nullhomotopic simple loop in \mathfrak{S} . Note that a point of the Teichmüller space yields a monodromy representation $\rho : \pi_1(\mathfrak{S}) \rightarrow \text{PSL}(2, \mathbb{R})$, defined up to conjugations in $\text{PSL}(2, \mathbb{R})$. The function $\mathbb{I}(\gamma) := |\text{trace}(\rho(\gamma))|$ is universally Laurent on $\mathcal{X}^+(\mathfrak{S})$; in fact, it is universally Laurent in the square-roots of the exponentiated shear coordinates $Z_e := \sqrt{X_e}$. It is known [FG06] [S20a] [GHK15] that any element of $\mathbf{L}(\mathcal{X}^+(\mathfrak{S}))$ is generated by these basic functions, together with relatively easy extra functions associated to punctures. So the problem of quantization almost boils down to quantizing these trace-of-monodromy functions $\mathbb{I}(\gamma)$.

The square-roots of the exponentiated shear coordinates appearing in $\mathbb{I}(\gamma)$ force us to replace the target space of the map in (1.1) by the bigger space $\mathcal{T}_\Delta^\omega$, which is the \mathbb{C} -algebra generated by the symbols \widehat{Z}_e and their inverses, mod out by the relations $\widehat{Z}_e \widehat{Z}_f = \omega^{2\epsilon_{ef}} \widehat{Z}_f \widehat{Z}_e$. Here $q = \omega^4$, and \mathcal{T}_Δ^q embeds into $\mathcal{T}_\Delta^\omega$ by $\widehat{X}_e \mapsto \widehat{Z}_e^2, \forall e \in T$.

The Bonahon-Wong *quantum trace* [BW11] is an algebra homomorphism

$$(1.2) \quad \text{Tr}_\Delta^\omega : \mathcal{S}^A(\mathfrak{S}) \rightarrow \mathcal{T}_\Delta^\omega, \quad \text{with } A = \omega^{-2}.$$

Here $\mathcal{S}^A(\mathfrak{S})$ is the *skein algebra* [P99, T91] of \mathfrak{S} at quantum parameter A , which is the \mathbb{C} -module generated by isotopy classes of framed unoriented links in the thickened surface $\mathfrak{S} \times (-1, 1)$ subject to certain local relations. The product of two framed links is given by stacking one above the other. The skein algebra $\mathcal{S}^A(\mathfrak{S})$ is known to be a quantization of the SL_2 -character variety of \mathfrak{S} in the direction of the Weil-Petersson-Goldman Poisson bracket (see [PS00] and references therein). Actually the image of the quantum trace map (1.2) is in a smaller subalgebra $\mathcal{Z}_\Delta^\omega$ of $\mathcal{T}_\Delta^\omega$, the skew-field of fractions of which the quantum coordinate change isomorphism of the Chekhov-Fock algebras can be extended to [H10] [BW11], and the quantum trace map is compatible with this extended quantum coordinate change. We will recall the definitions of the skein algebra, the Bonahon-Wong quantum trace map, as well as all these facts in Section 3.

Any simple loop γ in \mathfrak{S} can be considered as a framed link in the thickened surface $\mathfrak{S} \times (-1, 1)$, using the standard blackboard framing. In Allegretti-Kim [AK17], the function $\mathbb{I}(\gamma)$ is quantized by setting

$$\mathbb{I}_\Delta^q(\gamma) = \text{Tr}_\Delta^\omega(\gamma).$$

We note that a full description of \mathbb{I}_Δ^q involves still more non-trivial ideas; see [AK17]. For example, the above recipe $\mathbb{I}_\Delta^q(\gamma) = \text{Tr}_\Delta^\omega(\gamma)$ does not immediately fit into the original setting as in eq.(1.1), but only into a certain square-root version. We also note that the Bonahon-Wong quantum trace enjoys some more properties in accordance with the conjecture of Fock-Goncharov [FG06] for \mathbb{I}_Δ^q ; see [AK17], and also [CKKO20] for an important positivity property.

1.3. Another quantization using Gabella quantum holonomy. Gabella [G17] constructed for every oriented simple loop γ in \mathfrak{S} a *quantum holonomy* $\text{TrHol}_\Delta^\omega(\gamma) \in \mathcal{T}_\Delta^\omega$. Gabella's construction works for general Lie groups SL_N , where the algebra $\mathcal{T}_\Delta^\omega$ is replaced by the Fock-Goncharov quantum moduli space. The construction is based on works of Gaiotto, Moore, Neitzke, Galakhov, Longhi, and others [G12] [GMN12] [GMN13] [GMN14] [GLM15], which give a correspondence between the GL_N holonomies for the surface \mathfrak{S} and the abelian GL_1 holonomies for certain N -fold branched cover of \mathfrak{S} . Building on these works, and combining with the idea of Bonahon-Wong about going to three dimensions, Gabella constructed the quantum holonomy $\text{TrHol}_\Delta^\omega(\gamma)$. In this paper we focus only on the case $N = 2$.

Let us briefly and informally discuss the main ingredients in Gabella's construction. Details will be given in Section 4. Let $\pi : \widetilde{\mathfrak{S}}_\Delta \rightarrow \mathfrak{S}$ be a branched double covering of \mathfrak{S} branched over a set $\mathcal{V} \subset \mathfrak{S}$ consisting of exactly one point in the interior of each triangle of the triangulation Δ . Let $\mathfrak{S}' = \mathfrak{S} \setminus \mathcal{V}$. Let $\gamma : [0, 1] \rightarrow \mathfrak{S}$ be a not-necessarily-closed smooth oriented curve in a general position. When the image of γ lies in \mathfrak{S}' , there are exactly two lifts of γ in $\widetilde{\mathfrak{S}}_\Delta$. If γ is considered up to homotopy in \mathfrak{S} fixing its endpoints, then it has many more mutually non-homotopic lifts in $\widetilde{\mathfrak{S}}_\Delta$. In Gaiotto-Moore-Neitzke's construction, based on the triangulation one chooses a certain finite number of lifts, called *admissible lifts*, of the homotopy class of γ in \mathfrak{S} . Then, the GL_2 parallel transport for γ is expressed as sum over all admissible lifts $\widetilde{\gamma}$ of a formal variable $X_{\widetilde{\gamma}}$ times a certain coefficient coming from physics, counting the so-called framed BPS states. Galakhov, Longhi and Moore proposed a quantum parallel transport

$$\sum_{\widetilde{\gamma}: \text{admissible lift}} q^{-\text{wr}(\widetilde{\gamma})} \widehat{X}_{\widetilde{\gamma}},$$

where $\text{wr}(\widetilde{\gamma}) \in \mathbb{Z}$ is the *writhe* of the lift $\widetilde{\gamma}$, i.e. the number of self-intersections of $\widetilde{\gamma}$ counted with signs, and $\widehat{X}_{\widetilde{\gamma}}$ is a certain formal variable associated to $\widetilde{\gamma}$. This construction works well for open curves γ . When γ is a closed oriented curve in \mathfrak{S} , one can represent it by a map $\gamma : [0, 1] \rightarrow \mathfrak{S}$ by choosing a basepoint, then apply the above construction, where the variables $\widehat{X}_{\widetilde{\gamma}}$ become monomials in $\mathcal{T}_\Delta^\omega$; in general, the result depends on the choice of basepoint. Gabella suggested a modification in order to remove this dependency, and hence to construct a quantum holonomy of a simple closed oriented curve γ in \mathfrak{S} . Assume γ is in general position with respect to the arcs of Δ . Choose an arc e of Δ and cut γ

along e to get a collection of open curves γ_i . Apply the Galakhov-Longhi-Moore construction to each of γ_i . The intersection of γ and a small neighborhood of e is then interpreted as a certain type of tangle in the thickening of this neighborhood, which can be viewed as a *biangle*, and a modification of the operator invariant à la Reshetikhin-Turaev [RT90] [T16] of this tangle is used to “glue” the quantum holonomy of the arcs γ_i to give the Gabella quantum holonomy $\text{TrHol}_\Delta^\omega(\gamma) \in \mathcal{T}_\Delta^\omega$ [G17]. Gabella showed that $\text{TrHol}_\Delta^\omega(\gamma)$ is an isotopy invariant of simple closed oriented curves γ in \mathfrak{S} . It should be noted that while the Bonahon-Wong quantum trace map is defined for non-oriented framed links in the thickening of \mathfrak{S} , the Gabella quantum holonomy is defined for oriented simple loop in \mathfrak{S} . The definition of the Gabella quantum holonomy, where lifting of curves and writhe numbers are used, is very different from that of the Bonahon-Wong quantum trace.

A special case of the main theorem of the present paper is the following.

Theorem 1.1 (Part of Theorem 5.2). *Suppose \mathfrak{S} is a punctured surface with an ideal triangulation Δ . Let γ be an oriented simple closed curve in \mathfrak{S} . The Bonahon-Wong quantum trace and the Gabella quantum holonomy for γ coincide with each other*

$$\text{Tr}_\Delta^\omega(\gamma) = \text{TrHol}_\Delta^\omega(\gamma).$$

Here in the left hand side we consider γ as an unoriented link with the blackboard framing.

In particular, the deformation quantization map \mathbb{I}_Δ^q of Allegretti-Kim using Bonahon-Wong quantum trace coincides with Gabella’s solution (after a certain modification to be explained), for the trace-of-monodromy function $\mathbb{I}(\gamma) \in \mathbf{L}(\mathcal{X}^+(\mathfrak{S}))$ of a non-nullhomotopic simple loop γ in \mathfrak{S} .

1.4. More general result, and proof. Actually we will prove a much more general result. First we show that one can extend the Gabella quantum homonomy to not only all framed oriented links in the thickening of \mathfrak{S} , but also to a larger class of so-called *stated VH-tangles* in surfaces with boundary, and for VH-tangles we show that the Bonahon-Wong quantum trace and the Gabella quantum homonomy are not exactly equal but related by a certain relation. The more general result for VH-tangles is actually needed for the proof, as we decompose the surface into triangles and biangles, and establish the relation in each biangle and each triangle. Then we show that we can “glue” the relations so obtained in the biangles and triangles to get the desired global relation.

First we allow the surface \mathfrak{S} to have boundary. So $\mathfrak{S} = \Sigma \setminus \mathcal{P}$, where Σ is a compact oriented surface with (possibly empty) boundary and \mathcal{P} is a finite set that intersects every connected component of the boundary of Σ . By a *tangle* K in the thickened surface $\mathfrak{S} \times (-1, 1)$ we mean an unoriented compact properly embedded framed 1-dimensional submanifold of $\mathfrak{S} \times (-1, 1)$. A *V-tangle* is a tangle K such that for each boundary arc b of \mathfrak{S} , the endpoints of K in $b \times (-1, 1)$ have distinct vertical coordinates, i.e. distinct elevations $\in (-1, 1)$. A *VH-tangle* is a V-tangle K such that for each boundary arc b , the endpoints of K in $b \times (-1, 1)$ have distinct horizontal coordinates, i.e. project down to distinct points in b . By *V-isotopy* and *VH-isotopy* we mean the isotopies within the respective classes. A *state* of a V-tangle or VH-tangle is a map from its boundary points to the set $\{+, -\}$. Bonahon and Wong [BW11] extend the definition of the skein algebra to the stated skein algebra $\mathcal{S}_s^A(\mathfrak{S})$ based on stated V-tangles, and show that the quantum trace map $\text{Tr}_\Delta^\omega : \mathcal{S}_s^A(\mathfrak{S}) \rightarrow \mathcal{T}_\Delta^\omega$ can be defined as an algebra homomorphism.

In Gabella’s definition the writhe plays an important role, but the V-isotopy can change the writhe. This is the reason why we have to consider the finer class of VH-tangles, as VH-isotopy does not change the writhe. For each VH-isotopy class of a stated VH-tangle α in $\mathfrak{S} \times (-1, 1)$ we will define the Gabella quantum holonomy $\text{TrHol}_\Delta^\omega(\alpha) \in \mathcal{T}_\Delta^\omega$, extending Gabella’s definition for simple closed curves in \mathfrak{S} .

In the process we reformulate Gabella’s construction of [G17] so that it is more easily understandable to mathematics community, provide a complete treatment of the biangle part and give precise interpretation in terms of the Reshetikhin-Turaev invariant, and use different normalization of the monomial part than Gabella’s. Various properties, including the well-definedness, were not completely proved in [G17], and will follow as a consequence of the present paper.

If we forget the H-structure in a stated VH-tangle α , we get a stated V-tangle. There is a simple quantity $\partial\mathcal{C}_\mathfrak{S}(\alpha) \in \mathbb{Z}$ (Def.5.1) which records some information from this forgetting process. The main result of the present paper is the following.

Theorem 1.2 (main theorem). *For a stated VH-tangle α in the thickened surface $\mathfrak{S} \times (-1, 1)$,*

$$\mathrm{Tr}_{\Delta}^{\omega}(\alpha) = \omega^{2\mathrm{wr}(\alpha)} \omega^{\partial\mathcal{C}_{\mathfrak{S}}(\alpha)} \mathrm{TrHol}_{\Delta}^{\omega}(\alpha),$$

where $\mathrm{wr}(\alpha)$ is the usual writhe of the tangle α in $\mathfrak{S} \times (-1, 1)$.

We briefly explain the idea of proof. For a biangle, we start from the well-known Reshetikhin-Turaev operator invariant [RT90] [T16], and show that a modification by a certain boundary twist and writhe still yields an operator invariant. We then show that the matrix elements of these operator invariants are precisely the Bonahon-Wong biangle quantum trace and Gabella biangle quantum holonomy. For a triangle, we first write the Bonahon-Wong's triangle factor as the so-called Weyl-ordered monomial times some power of ω which we call a *deviation*. Our main technical lemma (Lem.5.9) is the formula expressing this deviation in terms of certain combination of writhe in \mathfrak{S} , writhe of (Gabella) lifts in $\tilde{\mathfrak{S}}_{\Delta}$, and a sign correction at the sides of the triangle. In the entire surface, we show that the boundary twists of biangles and the sign corrections of triangles cancel each other at inner arcs of Δ , whereas the two kinds of writhes add up throughout the surface.

While the twist factor $\omega^{2\mathrm{wr}(\alpha)} \omega^{\partial\mathcal{C}_{\mathfrak{S}}(\alpha)}$ is indeed what makes possible the forgetting process from VH-tangles to V-tangles, it also records the orientation information of the tangle α . In particular, even if α is a closed tangle, so that it has no V-, H-, VH-structure nor a state, the Bonahon-Wong quantum trace and the Gabella quantum holonomy are not exactly the same, but are related by $\mathrm{Tr}_{\Delta}^{\omega}(\alpha) = \omega^{2\mathrm{wr}(\alpha)} \mathrm{TrHol}_{\Delta}^{\omega}(\alpha)$.

Acknowledgments. T.T.Q. Lê is partially supported by the National Science Foundation under Grant No. DMS 1811114. For H. Kim: This research was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education(grant number 2017R1D1A1B03030230). This work was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIT) (No. 2020R1C1C1A01011151). H. Kim thanks Dylan Allegretti, Carlos Scarinci, Linhui Shen, Maxime Gabella and Francis Bonahon for help, discussion, questions, and comments.

2. QUANTUM TEICHMÜLLER THEORY

We establish basic definitions for set-up, and briefly introduce quantum Teichmüller theory.

2.1. Ideal triangulation of a non-compact surface. In this subsection we choose to mostly follow the terminology convention of [L17], with some modification.

Definition 2.1 ([L17]). • *A generalized marked surface is a pair (Σ, \mathcal{P}) , where Σ is a compact oriented surface with boundary $\partial\Sigma$ and \mathcal{P} is a finite subset of Σ , such that each component of $\partial\Sigma$ should have at least one point of \mathcal{P} .*

• *The surface orientation on Σ determines the notion of clockwise/counterclockwise rotation on (the tangent space of each point of) the surface, and the convention of the boundary-orientation on the boundary $\partial\Sigma$ coming from the surface orientation is chosen to be compatible with the clockwise rotation on (points of) surface near boundary.*

• *Elements of \mathcal{P} are called marked points. Elements of \mathcal{P} in $\Sigma \setminus \partial\Sigma$ are called interior marked points, or punctures.*

• *If $\partial\Sigma = \emptyset$, then (Σ, \mathcal{P}) is called a punctured surface.*

• *By the boundary of the generalized marked surface (Σ, \mathcal{P}) we mean $(\partial\Sigma) \setminus \mathcal{P}$, and write as $\partial\Sigma \setminus \mathcal{P}$.*

• *A diffeomorphism between generalized marked surfaces (Σ, \mathcal{P}) and (Σ', \mathcal{P}') is an orientation-preserving diffeomorphism $\Sigma \rightarrow \Sigma'$ that restricts to a bijection $\mathcal{P} \rightarrow \mathcal{P}'$. We say (Σ, \mathcal{P}) and (Σ', \mathcal{P}') are diffeomorphic if there exists a diffeomorphism between them.*

As mentioned in [L17], a generalized marked surface defined this way corresponds to a ‘punctured surface with boundary’ of [BW11].

Definition 2.2. *By the phrase “with boundary” we always mean “with (possibly empty) boundary”.*

In particular, in Def.2.1, we allow $\partial\Sigma$ to be empty. Notice that, in Def.2.1, in case $\partial\Sigma$ is empty, the boundary $\partial\Sigma \setminus \mathcal{P}$ of (Σ, \mathcal{P}) is also empty.

Definition 2.3 ([L17]). *Let (Σ, \mathcal{P}) be a generalized marked surface.*

• A \mathcal{P} -link is an immersion $\alpha : C \rightarrow \Sigma$, where C is a compact non-oriented 1-manifold with boundary, such that

- (1) restriction of α onto the interior of C is an embedding into $\Sigma \setminus \mathcal{P}$, and
- (2) α maps the boundary of C into \mathcal{P} .

The interior of this \mathcal{P} -link is the image under α of the interior of C , and is denoted by $\mathring{\alpha}$.

- When C is $[0, 1]$, we call α a \mathcal{P} -arc. When C is S^1 , we call α a \mathcal{P} -knot.
- When C has no boundary, we call α a closed \mathcal{P} -link.
- Two \mathcal{P} -links are \mathcal{P} -isotopic if they are isotopic in the class of \mathcal{P} -links.
- A \mathcal{P} -arc is called a boundary arc if it is contained in $\partial\Sigma$.
- A \mathcal{P} -link is often identified with its image in Σ .

In the literature, the surface is commonly considered to be $\Sigma \setminus \mathcal{P}$, in which case a \mathcal{P} -arc corresponds to a so-called *ideal arc*, a \mathcal{P} -knot to a simple loop in $\Sigma \setminus \mathcal{P}$, and a closed \mathcal{P} -link to a simple closed curve in $\Sigma \setminus \mathcal{P}$. The last line of Def.2.3 indicates that a \mathcal{P} -link is an unoriented object. One easy observation is that the boundary $\partial\Sigma \setminus \mathcal{P}$ is the disjoint union of interiors of boundary arcs, which are the connected components of $\partial\Sigma \setminus \mathcal{P}$. The interior of a boundary arc may also be called a boundary arc, by abuse of notation. We now go on to triangulations; we follow conventions of [L17] [FST08].

Definition 2.4 (e.g. [L17] [FST08]). *A generalized marked surface (Σ, \mathcal{P}) is triangulable if and only if \mathcal{P} is non-empty and (Σ, \mathcal{P}) is not diffeomorphic to one of the following:*

- a sphere (with no boundary) with one or two marked points
- a monogon with no interior marked point, (i.e. genus 0, with one boundary component, with one marked point on the boundary, and no interior marked point) or
- a bigon with no interior marked point (i.e. genus 0, with one boundary component, with two marked points on the boundary, and no interior marked point), also called a biangle. ■

We do not always assume that (Σ, \mathcal{P}) is triangulable, although we do when we talk about triangulations. Later, it is crucial that we shall also need to consider the third case of the above definition.

Definition and Lemma 2.5 (see e.g. [L17] [FST08]). *Let (Σ, \mathcal{P}) be a triangulable generalized marked surface.*

• For integer $n \geq 3$, a \mathcal{P} - n -gon is a smooth map $\beta : \sigma \rightarrow \Sigma$ from a regular n -gon σ in the standard plane \mathbb{R}^2 to Σ , such that

- (1) the restriction of β onto the interior $\mathring{\sigma}$ is a diffeomorphism onto its image, and
- (2) the restriction of β onto each of the n sides of σ is a \mathcal{P} -arc, called a side (or edge) of β .

In particular, a \mathcal{P} -3-gon is called a \mathcal{P} -triangle.

• If two sides of a \mathcal{P} - n -gon coincide as a \mathcal{P} -arc, such a side, as well as that \mathcal{P} - n -gon, is said to be self-folded.

• A \mathcal{P} - n -gon is oriented if the relevant map $\beta : \sigma \rightarrow \Sigma$ is orientation-preserving, where σ is oriented according to the standard orientation on \mathbb{R}^2 .

• A \mathcal{P} -triangulation (or, a triangulation, when \mathcal{P} is clear) of Σ (or, an ideal triangulation or just a triangulation of (Σ, \mathcal{P})) is a collection Δ of \mathcal{P} -arcs such that

- (1) no \mathcal{P} -arc in Δ bounds a disk whose interior is in $\Sigma \setminus \mathcal{P}$,
- (2) no two \mathcal{P} -arcs in Δ intersect in $\Sigma \setminus \mathcal{P}$ and no two are \mathcal{P} -isotopic, and
- (3) Δ is a maximal collection of \mathcal{P} -arcs with the above properties (1)–(2).

For a triangulation Δ , one can replace \mathcal{P} -arcs in Δ by \mathcal{P} -arcs in their respective \mathcal{P} -isotopy classes such that every arc in Δ isotopic to a boundary arc is a boundary arc. We always assume that Δ satisfies this condition.

• An element of Δ is called a (constituent) edge of the triangulation Δ . An edge of Δ is called a boundary edge of Δ if it is a boundary arc, and an internal edge of Δ otherwise. Let $\mathring{\Delta}$ be the set of all internal edges of Δ .

- For a \mathcal{P} -triangulation Δ , the closure of each connected component of $\Sigma \setminus (\bigcup_{e \in \Delta} e)$ can be naturally given a structure of an oriented \mathcal{P} -triangle. By a (constituent) (\mathcal{P} -)triangle of Δ we mean one of these triangles coming from Δ . Denote by $\mathcal{F}(\Delta)$ the set of all triangles of Δ .

- We say (Σ, \mathcal{P}) is triangulated if it is equipped with a \mathcal{P} -triangulation.

The notions of \mathcal{P} -triangulation and \mathcal{P} -triangle correspond to those of *ideal triangulation* and *ideal triangle* in the literature. Implicitly or explicitly, we will identify two triangulations if one can be obtained by simultaneous isotopy of (edges of) the other; when extra care is needed, we shall make it clear. For convenience of the present paper, we define the notion of corner of a triangle as follows.

Definition and Lemma 2.6. *Let t be a constituent triangle of a triangulation Δ of a generalized marked surface (Σ, \mathcal{P}) . Let $\beta : \sigma \rightarrow \Sigma$ be a map giving t an oriented \mathcal{P} -triangle structure. Let e, f, g be the sides of σ , appearing clockwise in this order. Denote by same labels e, f, g the corresponding sides of t . The three pairs of sides (e, f) , (f, g) , and (g, e) are called corners of t . The three corners of t are well-defined and mutually distinct.*

Visually, one can consider a corner (e, f) of t as being a small part of $t \setminus (e \cup f \cup g)$ close to the vertex of the t shared by e, f , lying in between e, f . There may arise some confusion when dealing with a self-folded triangle. In the above definition, we give three different labels e, f, g for the three sides; however, sometimes we identify a side with its image, in which case we will need only two distinct labels for the sides of a self-folded triangle, say e, e, f . Notice that, still in such a case, the three corners (e, e) , (e, f) , (f, e) of this triangle are unambiguously defined. However, notice that a pair of edges of Δ may not be sufficient to determine a corner, because sometimes such a pair may represent corners of two different triangles.

We find it convenient to define the following notion here.

Definition 2.7. *Let Δ be a triangulation of a generalized marked surface (Σ, \mathcal{P}) , and let γ be a closed \mathcal{P} -link in Σ . Denote by $\#(\gamma \cap \Delta)$ the total number of intersections of γ and the edges of Δ ; it can be infinite. We say that γ is in a minimal position with respect to Δ if $\#(\gamma \cap \Delta)$ is minimal among the numbers $\#(\gamma' \cap \Delta)$, where γ' runs through all \mathcal{P} -knots that are \mathcal{P} -isotopic to γ .*

It is easy to see that if γ is in a minimal position with respect to Δ , then $\#(\gamma \cap \Delta)$ is finite. It is well-known that the intersection numbers $\#(\gamma \cap e)$ with the edges $e \in \Delta$ completely determine the \mathcal{P} -knot γ in a minimal position, up to \mathcal{P} -isotopy.

Lemma 2.8. *Let Δ be a triangulation of a generalized marked surface (Σ, \mathcal{P}) , and let γ be a closed \mathcal{P} -link in Σ . Let γ' be a closed \mathcal{P} -link that is \mathcal{P} -isotopic to γ and is in a minimal position with respect to Δ . For each $e \in \Delta$, denote by $a_e(\gamma) \in \mathbb{Z}_{\geq 0}$ the number of intersections of γ' and e , called the intersection number of γ and e .*

The intersection numbers completely determine a closed \mathcal{P} -link γ up to \mathcal{P} -isotopy.

For a proof, we refer e.g. to [FG06]; the intersection numbers are 2 times the Fock-Goncharov tropical \mathcal{A} -coordinate of γ , and it is straightforward to re-construct γ out of these numbers.

As is widely used in the literature, counting of corners effectively encodes the combinatorics of Δ .

Definition 2.9. *Let Δ be a triangulation of a generalized marked surface (Σ, \mathcal{P}) . Let e, f be constituent edges of Δ , and let t be a constituent triangle of Δ . Define*

$$\begin{aligned} c_{ef}(t) &= c_{ef}^\Delta(t) := \begin{cases} 1 & \text{if } (e, f) \text{ is a corner of } t, \\ 0 & \text{otherwise,} \end{cases} \\ \varepsilon_{ef}(t) &= \varepsilon_{ef}^\Delta(t) := c_{ef}(t) - c_{fe}(t), \\ \varepsilon_{ef} &= \varepsilon_{ef}^\Delta := \sum_{t \in \mathcal{F}(\Delta)} \varepsilon_{ef}(t). \end{aligned}$$

The matrix $\varepsilon = \varepsilon^\Delta = (\varepsilon_{ef})_{e, f \in \Delta}$ is called the exchange matrix of Δ .

The matrix ε is sometimes called the signed adjacency matrix (e.g. [FST08]), or a face matrix (e.g. [L17]). Notice that the matrix $B = (b_{ij})$ of [FST08] coincides with the above $\varepsilon = (\varepsilon_{ij})$ in case Δ has no self-folded triangles, and otherwise it differs from ε in general.

2.2. Teichmüller space and shear coordinates. The basic geometric objects of study are certain versions of the so-called Teichmüller space of a surface (Σ, \mathcal{P}) . In general, the Teichmüller space of a surface refers to the space of all complete hyperbolic metrics on the surface, considered up to pullback by diffeomorphisms isotopic to the identity. In case of a generalized marked surface, some care is needed for the behavior near the boundary and marked points. The most general case appears in Dylan Allegretti's thesis [A16], which we follow. Like in [A16], we formulate using the monodromy representations, instead of directly dealing with hyperbolic metrics.

Definition 2.10 (enhanced Teichmüller space of punctured surface; see e.g. [FG06]). *Let (Σ, \mathcal{P}) be a triangulable generalized marked surface, and suppose that (Σ, \mathcal{P}) is a punctured surface, i.e. $\partial\Sigma = \emptyset$.*

- The Teichmüller space $\mathcal{T}(\Sigma, \mathcal{P})$ is the set

$$(2.1) \quad \text{Hom}^{\text{df}}(\pi_1(\Sigma \setminus \mathcal{P}), \text{PSL}(2, \mathbb{R})) / \text{PSL}(2, \mathbb{R}).$$

More precisely, $\mathcal{T}(\Sigma, \mathcal{P})$ consists of all group homomorphisms from $\pi_1(\Sigma \setminus \mathcal{P})$ to $\text{PSL}(2, \mathbb{R})$ that are faithful and have discrete image, where an element of $\mathcal{T}(\Sigma, \mathcal{P})$ is defined up to conjugation by an element of $\text{PSL}(2, \mathbb{R})$.

- Given a point of $\mathcal{T}(\Sigma, \mathcal{P})$, i.e. an equivalence class of a group homomorphism $\rho : \pi_1(\Sigma \setminus \mathcal{P}) \rightarrow \text{PSL}(2, \mathbb{R})$, an element p of \mathcal{P} is called a cusps (with respect to ρ) if the image under ρ of a small loop surrounding p is a parabolic element of $\text{PSL}(2, \mathbb{R})$, and called a hole (with respect to ρ) if this image is a hyperbolic element of $\text{PSL}(2, \mathbb{R})$.

- The enhanced Teichmüller space $\mathcal{X}^+(\Sigma, \mathcal{P})$ is the set of all pairs (ρ, O) , where ρ is a point of $\mathcal{T}(\Sigma, \mathcal{P})$, and O is the choice of an orientation for (a loop surrounding) each hole with respect to ρ .

Definition 2.11 (enhanced Teichmüller space for a triangulable generalized marked surface; [A16]). *Let (Σ, \mathcal{P}) be a triangulable generalized marked surface with nonempty boundary $\partial\Sigma \neq \emptyset$.*

- Let $(\Sigma^{\text{op}}, \mathcal{P}^{\text{op}})$ be the generalized marked surface same as (Σ, \mathcal{P}) , except equipped with the opposite orientation. Choose a parametrization for each boundary component of Σ , which induces that for the corresponding boundary component of Σ^{op} . Glue Σ and Σ^{op} along these parametrized boundary components, to construct a smooth oriented surface $\Sigma_{\mathbb{D}}$ without boundary; denote by $\mathcal{P}_{\mathbb{D}}$ the marked points on $\Sigma_{\mathbb{D}}$ coming from \mathcal{P} and \mathcal{P}^{op} . The resulting generalized marked surface $(\Sigma_{\mathbb{D}}, \mathcal{P}_{\mathbb{D}}) =: (\Sigma, \mathcal{P})_{\mathbb{D}}$, which is a punctured surface, is called the doubled surface for (Σ, \mathcal{P}) . Let $\iota : \Sigma_{\mathbb{D}} \rightarrow \Sigma_{\mathbb{D}}$ be the natural involutive diffeomorphism exchanging Σ and Σ^{op} , induced by the identification $\Sigma \leftrightarrow \Sigma^{\text{op}}$.

- The (generalized) enhanced Teichmüller space $\mathcal{X}^+(\Sigma, \mathcal{P})$ is defined as the ι -invariant subspace of the enhanced Teichmüller space $\mathcal{X}^+(\Sigma_{\mathbb{D}}, \mathcal{P}_{\mathbb{D}})$ of the doubled surface $(\Sigma_{\mathbb{D}}, \mathcal{P}_{\mathbb{D}})$, where $\mathcal{X}^+(\Sigma_{\mathbb{D}}, \mathcal{P}_{\mathbb{D}})$ is defined as in Def.2.10, on which ι acts naturally.

A starting point of many problems related to Teichmüller spaces is a construction of suitable coordinate systems for $\mathcal{X}^+(\Sigma, \mathcal{P})$. Among various kinds, what is relevant to the current situation is Thurston's shear coordinate function [T80], which makes use of the choice of an ideal triangulation.

Proposition 2.12 (Thurston-Fock's theorem, see e.g. [P12] [A16]). *Let Δ be a triangulation of a triangulable generalized marked surface (Σ, \mathcal{P}) . There exists a global coordinate system for the enhanced Teichmüller space $\mathcal{X}^+(\Sigma, \mathcal{P})$, whose coordinate functions are enumerated by the internal edges e of the triangulation Δ . These coordinate functions $X_e = X_e^{\Delta} : \mathcal{X}^+(\Sigma, \mathcal{P}) \rightarrow \mathbb{R}_{>0}$ associated to $e \in \mathring{\Delta}$, each of which is called the exponentiated shear coordinate for the internal edge $e \in \Delta$, provide a bijection*

$$\mathcal{X}^+(\Sigma, \mathcal{P}) \rightarrow (\mathbb{R}_{>0})^{\mathring{\Delta}} : (\rho, O) \mapsto X_e(\rho, O).$$

For geometric meaning of these coordinate functions, see e.g. [A16] or [P12]. Roughly speaking, in terms of hyperbolic metrics, they are the 'shearing' amounts that measure how the hyperbolic ideal triangles are glued along their edges to one another, to form the (hyperbolic) surface $\Sigma \setminus \mathcal{P}$. A key point of studying these coordinates is how they transform under change of ideal triangulations.

Proposition 2.13. *Let Δ and Δ' be ideal triangulations of a triangulable generalized marked surface (Σ, \mathcal{P}) . Then the exponentiated shear coordinate functions X'_e 's for the internal edges e' of Δ' can be expressed as rational functions in terms of those X_e 's for the internal edges e of Δ .*

2.3. Quantum Teichmüller space. We now review quantum Teichmüller space, which was first constructed in 1990's by Kashaev [K98] and independently by Chekhov-Fock [CF99] [F97]. For our purposes we follow the latter works, and also their modern developments as appearing e.g. in [L09] [BW11] [H10] [FG09].

For each triangulation Δ , we considered the exponentiated shear coordinate functions X_e associated to internal edges $e \in \mathring{\Delta}$. They provide identification of the (generalized) enhanced Teichmüller space $\mathcal{X}^+(\Sigma, \mathcal{P})$ with $(\mathbb{R}_{>0})^{\mathring{\Delta}}$, making $\mathcal{X}^+(\Sigma, \mathcal{P})$ a smooth manifold. This manifold $\mathcal{X}^+(\Sigma, \mathcal{P})$ is equipped with a natural Poisson structure, named Weil-Petersson Poisson structure, whose Poisson brackets among the coordinate functions are given by

$$\{X_e, X_f\} = \varepsilon_{ef} X_e X_f.$$

Thus one can verify that $\{X_e^{\pm 1} : e \in \mathring{\Delta}\}$ generates a Poisson subalgebra of the ring $C^\infty(\mathcal{X}^+(\Sigma, \mathcal{P}))$ of smooth functions on $\mathcal{X}^+(\Sigma, \mathcal{P})$. The following non-commutative algebras serve as quantum deformed versions of this Poisson subalgebra.

Definition 2.14 ([L09]). *Let Δ be a triangulation of a generalized marked surface (Σ, \mathcal{P}) , and $\mathring{\Delta}$ be the set of all internal edges of Δ . Let $\hbar \in \mathbb{R}$ be a quantum parameter, and let $q := \exp(\pi i \hbar) \in \mathbb{C}^*$.*

The internal Chekhov-Fock algebra $\mathring{\mathcal{T}}_\Delta^q$ is an algebra over \mathbb{C} defined by generators and relations:

$$\begin{aligned} \text{generating set} &: \{ \widehat{X}_e, \widehat{X}_e^{-1} \mid e \in \mathring{\Delta} \}, \\ \text{relations} &: \widehat{X}_e \widehat{X}_f = q^{2\varepsilon_{ef}} \widehat{X}_f \widehat{X}_e, \quad \forall e, f \in \mathring{\Delta}, \quad \widehat{X}_e \widehat{X}_e^{-1} = \widehat{X}_e^{-1} \widehat{X}_e = 1, \quad \forall e \in \mathring{\Delta}. \end{aligned}$$

The Chekhov-Fock algebra \mathcal{T}_Δ^q is an algebra over \mathbb{C} defined by generators and relations as follows:

$$\begin{aligned} \text{generating set} &: \{ \widehat{X}_e, \widehat{X}_e^{-1} \mid e \in \Delta \}, \\ \text{relations} &: \widehat{X}_e \widehat{X}_f = q^{2\varepsilon_{ef}} \widehat{X}_f \widehat{X}_e, \quad \forall e, f \in \Delta, \quad \widehat{X}_e \widehat{X}_e^{-1} = \widehat{X}_e^{-1} \widehat{X}_e = 1, \quad \forall e \in \Delta. \end{aligned}$$

Notice that $\mathring{\mathcal{T}}_\Delta^q$ and \mathcal{T}_Δ^q coincide when (Σ, \mathcal{P}) is a punctured surface, which is often the case in the literature. As the generators of the classical algebra are enumerated by internal edges, of course $\mathring{\mathcal{T}}_\Delta^q$ is more natural to consider than \mathcal{T}_Δ^q , but we shall also need \mathcal{T}_Δ^q later. As noted e.g. in [L09] [BW11] [H10], these algebras $\mathring{\mathcal{T}}_\Delta^q$ and \mathcal{T}_Δ^q satisfy the so-called Ore Condition of ring theory [BG02] [C95], and therefore their skew-fields (i.e. division algebras) of fractions $\text{Frac}(\mathring{\mathcal{T}}_\Delta^q)$ and $\text{Frac}(\mathcal{T}_\Delta^q)$ can be considered. As written in [L09], elements of $\text{Frac}(\mathcal{T}_\Delta^q)$ are formal fraction expressions PQ^{-1} , with $P, Q \in \mathcal{T}_\Delta^q$, $Q \neq 0$, where two such expressions $P_1Q_1^{-1}$ and $P_2Q_2^{-1}$ represent the same element of $\text{Frac}(\mathcal{T}_\Delta^q)$ if there exist nonzero $S_1, S_2 \in \mathcal{T}_\Delta^q$ such that $P_1S_1 = P_2S_2$ and $Q_1S_1 = Q_2S_2$. Product of such expressions can also be written as one fraction PQ^{-1} by algebraic manipulation. Similarly for $\text{Frac}(\mathring{\mathcal{T}}_\Delta^q)$.

Proposition 2.15 ([CF99] [K98] [L09]). *There exists a skew-field isomorphism*

$$\Phi_{\Delta, \Delta'}^q : \text{Frac}(\mathcal{T}_{\Delta'}^q) \rightarrow \text{Frac}(\mathcal{T}_\Delta^q)$$

associated to each pair Δ, Δ' of triangulations of a generalized marked surface (Σ, \mathcal{P}) , satisfying:

- (1) When $q = 1$ it recovers the coordinate change formulas for exponentiated shear coordinates for internal edges,
- (2) The consistency relation holds

$$\Phi_{\Delta, \Delta''}^q = \Phi_{\Delta, \Delta'}^q \circ \Phi_{\Delta', \Delta''}^q$$

for each triple $\Delta, \Delta', \Delta''$ of triangulations.

In fact, the above proposition was first established for the internal Chekhov-Fock algebras only, i.e. for the maps $\text{Frac}(\mathring{\mathcal{T}}_{\Delta'}^q) \rightarrow \text{Frac}(\mathring{\mathcal{T}}_\Delta^q)$. We note that these original results naturally extend to the Chekhov-Fock algebras as written above.

Some authors (e.g. [L09] [BW11]) use the word quantum Teichmüller space as referring to the quotient of disjoint union of all $\text{Frac}(\mathcal{T}_\Delta^q)$ by the equivalence relation given by the identifications $\Phi_{\Delta, \Delta'}^q$. We might also be doing so from time to time, but we do not make a serious use of this word.

2.4. Deformation quantization of Teichmüller space and the quantum ordering problem.

The isomorphisms in Prop.2.15 let us identify $\text{Frac}(\mathcal{T}_\Delta^q)$ for different Δ 's in a consistent way. However, it is $\text{Frac}(\mathcal{T}_\Delta^q)$ that is being identified with others, instead of the original Chekhov-Fock algebra \mathcal{T}_Δ^q which we began with. For certain reasons, among elements of $\text{Frac}(\mathcal{T}_\Delta^q)$, those belonging to \mathcal{T}_Δ^q are considered more important. First, in order to be physically relevant, one must realize elements of the quantum algebra as operators on Hilbert space. It is relatively easy to deal with representation of the algebra \mathcal{T}_Δ^q on the Hilbert space using functional analysis, but hard to do for its skew-field of fractions. Second reason comes from the viewpoint of considering the enhanced Teichmüller space as a cluster \mathcal{X} -variety. Each triangulation yields a chart, and regular functions on this chart are defined as Laurent polynomials in the exponentiated shear coordinates; we note that Laurent polynomials play a crucial role in the theory of cluster algebras and cluster varieties. Quantum regular functions for the quantized chart are non-commutative Laurent polynomials in the quantum counterpart of the exponentiated shear coordinates.

Hence, for both reasons coming from representation theory and cluster variety theory, the nice functions to deal with, for classical and quantum as well, are functions that are regular for every chart, i.e. that are Laurent polynomials in every triangulation.

Definition 2.16. *Define*

$$\mathbf{L}_\Delta^q := \bigcap_{\Delta'} \Phi_{\Delta, \Delta'}^q(\mathcal{T}_{\Delta'}^q) \subset \mathcal{T}_\Delta^q \subset \text{Frac}(\mathcal{T}_\Delta^q).$$

where the intersection runs through all ideal triangulations Δ' of the relevant generalized marked surface (Σ, \mathcal{P}) , including Δ . Elements of $\text{Frac}(\mathcal{T}_\Delta^q)$ that belong to \mathbf{L}_Δ^q are said to be universally Laurent.

Lemma 2.17. $\Phi_{\Delta, \Delta'}^q$ induces isomorphism from $\mathbf{L}_{\Delta'}^q$ to \mathbf{L}_Δ^q . ■

So the algebras \mathbf{L}_Δ^q for different Δ 's can be consistently identified, so they can be collectively denoted as $\mathbf{L}^q = \mathbf{L}_{(\Sigma, \mathcal{P})}^q$. That is, in the style used in [BW11], we can view $\mathbf{L}^q = \bigsqcup_{\Delta} \mathbf{L}_\Delta^q / \sim$, where \bigsqcup is the set disjoint union and \sim is the equivalence relation coming from $\Phi_{\Delta, \Delta'}^q$.

The discussion so far is only on construction of a consistent quantum system related to the classical system, which is the enhanced Teichmüller space with the Weil-Petersson Poisson structure. That is, we now have algebra of quantum observables, namely \mathbf{L}^q , independent of the choice of an ideal triangulation Δ . Next step is to establish a quantization, which is a map from the algebra of classical observables to the algebra of quantum observables. Namely, given a classical observable, i.e. a smooth function on the enhanced Teichmüller space $\mathcal{X}^+(\Sigma, \mathcal{P})$, what quantum observable do we assign to it? A place to begin with is each of the exponentiated shear coordinate functions X_e for a chosen ideal triangulation Δ . A natural candidate of a quantization map is to send each X_e to $\widehat{X}_e \in \mathcal{T}_\Delta^q$. Then, what about other functions? At the moment, let us only focus on the functions on $\mathcal{X}^+(\Sigma, \mathcal{P})$ that can be written as Laurent polynomials in X_e 's for a given Δ , such as $X_e X_f + X_g^{-2}$. To each such Laurent polynomial in X_e , we'd like to assign a non-commutative Laurent polynomial in \widehat{X}_e 's that recovers the classical one when we put $q = 1$ and replace each \widehat{X}_e by X_e . For a fixed Δ , building such an assignment so that it satisfies the axiom of a 'deformation quantization map' is not so hard. The difficult part is to make sure that such a deformation quantization map does not depend on the choice of Δ , in a sense.

The first step is to restrict our attention to (classical) functions on $\mathcal{X}^+(\Sigma, \mathcal{P})$ that can be written as Laurent polynomials in the exponentiated shear coordinates, for *every* ideal triangulations; finding all of them is already a highly nontrivial task, and is accomplished in [FG06]. Then, for each Δ , devise a way to assign to each such universally Laurent function a quantum Laurent polynomial, and prove that the resulting quantum Laurent polynomials for different Δ 's are related to each other by the quantum mutation maps $\Phi_{\Delta, \Delta'}^q$.

To give an intuition, let us consider a simple toy model. To a function $X_e X_f$, what should we assign? Options are $\widehat{X}_e \widehat{X}_f$, $\widehat{X}_f \widehat{X}_e$, $q^r \widehat{X}_e \widehat{X}_f$ for some $r \in \mathbb{Z}$, or maybe it could be even more complicated. Which one is the best choice? For a more general Laurent polynomial, we should choose how to quantize each monomial term. Finding a good choice of quantum Laurent polynomial so that it satisfies certain favorable properties is sometimes referred to as the quantum ordering problem, as if it is the problem of choosing the order of product of non-commutative quantum functions.

For a monomial, there is a well-known standard answer, namely the Weyl-ordered product; we formulate this in a general setting as follows.

Definition 2.18 (Weyl-ordered product). *Let $\mathbf{X}_1, \dots, \mathbf{X}_n$ be elements living in an algebra, and satisfies*

$$\mathbf{X}_i \mathbf{X}_j = A^{2m_{ij}} \mathbf{X}_j \mathbf{X}_i, \quad \forall i, j \in \{1, \dots, n\},$$

for some scalar A and integers $m_{ij} \in \mathbb{Z}$. In such a situation, we say that these n elements A -commute with one another. Define the Weyl-ordered product of these A -commuting elements as

$$[\mathbf{X}_1 \mathbf{X}_2 \cdots \mathbf{X}_n]_{\text{Weyl}} := A^{-\sum_{1 \leq i < j \leq n} m_{ij}} \mathbf{X}_1 \mathbf{X}_2 \cdots \mathbf{X}_n.$$

It is a straightforward exercise that the Weyl-ordered product is invariant under permutation, namely $[\mathbf{X}_1 \cdots \mathbf{X}_n]_{\text{Weyl}}$ equals $[\mathbf{X}_{\sigma(1)} \cdots \mathbf{X}_{\sigma(n)}]_{\text{Weyl}}$ for each permutation σ of $\{1, \dots, n\}$; it makes the Weyl-ordered product a standard answer to the quantum ordering problem of a monomial. However, for our case, for a classical function given by a Laurent polynomial, the quantum Laurent polynomial obtained by replacing each constituent monomial $X_e X_f \cdots$ by the Weyl-ordered product $[\widehat{X}_e \widehat{X}_f \cdots]_{\text{Weyl}}$ turns out *not* to satisfy the desired property, namely the compatibility under the quantum mutations $\Phi_{\Delta, \Delta'}^q$. So the quantum ordering problem for universally Laurent functions cannot be solved just by term-by-term Weyl-ordered products. In the present paper, we will review two solutions to this problem, namely one by Allegretti-Kim [AK17] and the other by Gabella [G17], and finally show that these two answers are the same.

3. BONAHO-WONG'S QUANTUM TRACE

3.1. Tangles and skein algebra. The known answers to the quantum ordering problem mentioned in the last section are heavily based on the work of Bonahon and Wong [BW11]. Bonahon-Wong's construction requires us to consider *tangles*, which are 1-dimensional submanifolds in a 3-dimensional manifold subject to certain conditions and with extra data. Depending on the authors and situations, these objects can be defined in different ways, and here we recollect some versions relevant to our paper.

Definition 3.1. • *For an oriented surface \mathfrak{S} with boundary, the thickening of \mathfrak{S} is the 3-manifold*

$$\mathfrak{S} \times (-1, 1),$$

whose boundary is $\partial(\mathfrak{S} \times (-1, 1)) = (\partial\mathfrak{S}) \times (-1, 1)$. We also say that $\mathfrak{S} \times (-1, 1)$ is a thickened surface.

- *The elevation of a point $(p, t) \in \mathfrak{S} \times (-1, 1)$ is the $(-1, 1)$ -coordinate, namely t .*
- *If A is a subset of \mathfrak{S} , we say that a point x of $\mathfrak{S} \times (-1, 1)$ lies over A if $x \in A \times (-1, 1)$.*

We will mostly apply this definition to

$$\mathfrak{S} = \Sigma \setminus \mathcal{P}$$

for a generalized marked surface (Σ, \mathcal{P}) . Note that connected components b of $\partial\mathfrak{S}$ are interiors of boundary arcs of (Σ, \mathcal{P}) , hence in particular are diffeomorphic to an open interval in \mathbb{R} ; as mentioned already, we refer to these b as boundary arcs of (Σ, \mathcal{P}) or of \mathfrak{S} . Note that the connected components of $\partial\mathfrak{S} \times (-1, 1)$, i.e. the boundary components of $\mathfrak{S} \times (-1, 1)$, are the thickenings $b \times (-1, 1)$ of boundary arcs b of \mathfrak{S} . The *projection* to \mathfrak{S} means the usual projection map $\mathfrak{S} \times (-1, 1) \rightarrow \mathfrak{S}$, or the image of some subset under this map.

Definition 3.2 (various tangles). *Let (Σ, \mathcal{P}) be a generalized marked surface, not necessarily triangulable. Let $\mathfrak{S} = \Sigma \setminus \mathcal{P}$.*

• *A tangle in a thickened surface $\mathfrak{S} \times (-1, 1)$ is a 1-dimensional compact manifold K with boundary properly embedded into $\mathfrak{S} \times (-1, 1)$ such that*

- (T1) *K is equipped with the choice of a framing on it, i.e. a continuous choice of a vector in $T_x(\mathfrak{S} \times (-1, 1)) \setminus T_x K$ for each point x of K , and*
- (T2) *the framing at each endpoint of K , i.e. a point of $\partial K \subset \partial\mathfrak{S} \times (-1, 1)$, is upward vertical, i.e. is parallel to the $(-1, 1)$ factor and points toward 1 of $(-1, 1)$.*
- *For a tangle K in $\mathfrak{S} \times (-1, 1)$, consider the conditions:*
 - (T3) *for each boundary arc b of \mathfrak{S} , the elevations of the endpoints of K lying over b are mutually distinct;*
 - (T4) *for each boundary arc b of \mathfrak{S} , no two endpoints of K lying over b project to a same point in b .*

The tangle K is called a V-tangle if it satisfies (T3), an H-tangle if it satisfies (T4), and a VH-tangle if it satisfies both (T3) and (T4).

- Define each of V-isotopy, H-isotopy, and VH-isotopy as an isotopy within the class of V-tangles, H-tangles, and VH-tangles, respectively.
- A V-tangle, H-tangle, or a VH-tangle is closed if the underlying tangle is closed.
- An oriented version of a tangle, V-tangle, H-tangle and VH-tangle, as well as the respective isotopies, can be defined.

In particular, by a tangle we always mean a framed tangle. In the three versions of tangles, the letter ‘V’ indicates that a V-isotopy class remembers the vertical ordering of endpoints lying over each boundary arc of \mathfrak{S} , while ‘H’ indicates that an H-isotopy class remembers the horizontal ordering of them. A tangle K in $\mathfrak{S} \times (-1, 1)$ can be represented by an embedding map $C \rightarrow \mathfrak{S} \times (-1, 1)$ of a compact 1-dimensional manifold C with boundary into $\mathfrak{S} \times (-1, 1)$. The tangles are often studied through their projections in the surface \mathfrak{S} , called *tangle diagrams*.

Definition 3.3. • A tangle diagram in an oriented surface \mathfrak{S} with boundary is a 1-dimensional compact manifold D with boundary properly immersed into \mathfrak{S} s.t. the interior of D lies in the interior of \mathfrak{S} , the boundary ∂D of D lies in the boundary $\partial \mathfrak{S}$, and the only possible singularities are transverse double self-intersections lying in the interior $\mathfrak{S} \setminus \partial \mathfrak{S}$ of \mathfrak{S} , together with the data:

- (TD1) for each self-intersection of D , called a crossing, the over- or under-passing information.
 In a picture, for each crossing x , the underpassing part of D is drawn as broken lines, i.e. a small neighborhood of x in this part deleted. The elements of ∂D are called the endpoints of D .

An isotopy of tangle diagrams is an isotopy within the class of tangle diagrams.

- A boundary-ordering on a tangle diagram D is the following data:
 (TD2) for each boundary arc b of \mathfrak{S} , the choice of a (total) ordering on the set $\partial_b D := \partial D \cap b$, called a vertical ordering on $\partial_b D$; if $x \in \partial_b D$ is higher than $y \in \partial_b D$ in this ordering, we write $x \succ y$.
- A tangle diagram D equipped with a boundary-ordering is called a boundary-ordered tangle diagram. When the boundary-ordering is clear from the context, it is denoted just by D .

An isotopy of boundary-ordered tangle diagrams is an isotopy within the class of boundary-ordered tangle diagrams.

- An oriented tangle diagram, a boundary-ordered oriented tangle diagram and their respective isotopies are defined analogously.

For a tangle K in $\mathfrak{S} \times (-1, 1)$ to yield a tangle diagram in \mathfrak{S} via projection, it must satisfy:

- (P1) the framing at every point of K is upward vertical, and
 (P2) the projection D of K via the projection $\mathfrak{S} \times (-1, 1) \rightarrow \mathfrak{S}$ is a tangle diagram.

It is known that each of a V-tangle, H-tangle, or a VH-tangle can be isotoped within the respective class to another one satisfying (P1) and (P2). For each case, the over- and under-passing information (TD1) for each crossing is naturally induced from K . For a V-tangle or a VH-tangle, the vertical ordering (TD2) on $\partial_b D$ for each boundary arc b of \mathfrak{S} is induced by the ordering on $\partial K \cap (b \times (-1, 1))$ coming from elevations; hence the name *vertical* ordering. We find it convenient to define also the notion of *horizontal* ordering on $\partial_b D$:

Definition 3.4. Let b be a boundary arc of a surface \mathfrak{S} (with boundary).

- The choice of an orientation on b induces an ordering on the points of b called the horizontal ordering on b as follows: a point $x \in b$ is (horizontally) higher than $y \in b$ if the direction from y to x matches the chosen orientation on b .
- Let D be a tangle diagram in \mathfrak{S} . The horizontal ordering on the set $\partial_b D$ with respect to an orientation on b is the ordering induced by the horizontal ordering on b .

In the present paper, if \mathfrak{S} is an oriented surface and no particular orientation on b is chosen, we use the boundary-orientation on b (Def.2.1) coming from the surface orientation on \mathfrak{S} .

Note that, once (P1) and (P2) are satisfied, a V-tangle or a VH-tangle in $\mathfrak{S} \times (-1, 1)$ yields a boundary-ordered tangle diagram in \mathfrak{S} via projection, while an H-tangle yields a tangle diagram. We say that

the respective diagram is a diagram of the respective tangle. Conversely, from an isotopy class of one of these three kinds of tangle diagrams one can recover a unique isotopy class of a respective kind of tangles in $\mathfrak{S} \times (-1, 1)$. However, non-isotopic tangle diagrams may yield isotopic tangles, if they differ by certain ‘moves’.

- (M1) framed Reidemeister moves I, II and III, as in Fig.1(I), (II) and (III); in each case, the left tangle and the right tangle differs only in a small disc where they differ as in the picture;
- (M2) boundary-exchange move as in Fig.1(IV), where the shaded region indicates the inside of the surface, the vertical thin line part of a boundary arc b , the left tangle D_1 and the right tangle D_2 differ only over a small half-disc near the boundary as in the picture, and the vertical orderings on $\partial_b D_1$ and $\partial_b D_2$ match each other with respect to the natural bijection $\partial_b D_1 \leftrightarrow \partial_b D_2$ sending x to x and y to y .




FIGURE 1. Elementary moves for tangles

Proposition 3.5 (correspondence between tangle diagrams and tangles; e.g. [O02]). *Let \mathfrak{S} be an oriented surface (with boundary).*

- (1) *Two tangle diagrams in \mathfrak{S} yield isotopic tangles in $\mathfrak{S} \times (-1, 1)$ if and only if they are related by a sequence of isotopies of tangle diagrams and moves (M1) and (M2).*
- (2) *Two tangle diagrams in \mathfrak{S} yield isotopic H-tangles in $\mathfrak{S} \times (-1, 1)$ if and only if they are related by a sequence of isotopies of tangle diagrams and moves (M1).*
- (3) *Two boundary-ordered tangle diagrams in \mathfrak{S} yield isotopic V-tangles in $\mathfrak{S} \times (-1, 1)$ if and only if they are related by a sequence of isotopies of tangle diagrams and moves (M1) and (M2).*
- (4) *Two boundary-ordered tangle diagrams in \mathfrak{S} yield isotopic VH-tangles in $\mathfrak{S} \times (-1, 1)$ if and only if they are related by a sequence of isotopies of tangle diagrams and moves (M1).*

The statements for oriented versions hold analogously.

We finally define the stated skein algebra through the following definitions.

Definition 3.6. *Let (Σ, \mathcal{P}) be a generalized marked surface, not necessarily triangulable. Let $\mathfrak{S} = \Sigma \setminus \mathcal{P}$.*

- *A state on a tangle K in $\mathfrak{S} \times (-1, 1)$ is an assignment of a sign to each endpoint of K , i.e. a map $s : \partial K \rightarrow \{+, -\}$. A stated tangle is a pair (K, s) of a tangle K and a state s on K . The stated versions of various kinds of tangles, such as stated V-tangles, are defined analogously.*

- *A state on a tangle diagram D is a map $s : \partial D \rightarrow \{+, -\}$. A stated tangle diagram is a pair (D, s) of a tangle diagram D and a state s on D . The stated versions of various kinds of tangle diagrams, such as stated boundary-ordered tangle diagrams, are defined analogously.*

- *If $\partial K = \emptyset$ or $\partial D = \emptyset$, the only possible state is denoted by \emptyset , and the stated tangle (K, \emptyset) and the stated tangle diagram (D, \emptyset) are said to be closed.*

Definition 3.7 (the skein algebra; [T91] [BW11] [L17] [L18]). *Let (Σ, \mathcal{P}) and \mathfrak{S} be as in Def.3.6.*

The (Kauffman bracket) stated skein algebra $\mathcal{S}_s^A(\Sigma, \mathcal{P})$ is the associative \mathbb{C} -algebra defined as follows.

- *The underlying \mathbb{C} -vector space is freely spanned by the isotopy classes of all possible stated V-tangles in $\mathfrak{S} \times (-1, 1)$, mod out by the following relations, where an element of $\mathcal{S}_s^A(\Sigma, \mathcal{P})$ represented by the stated tangle (K, s) is denoted by $[K, s]$, and is called a stated skein:*

- (SA1) *(Kauffman bracket skein relation) For each triple of stated V-tangles (K_1, s_1) , (K_0, s_0) and (K_∞, s_∞) that differ only over a small open disc in the interior of \mathfrak{S} as in Fig.2, one has:*

$$[K_1, s_1] = A^{-1}[K_0, s_0] + A[K_\infty, s_\infty]$$




FIGURE 2. Kauffman triple of tangles

(SA2) (the trivial loop relation) For each pair of stated V -tangles (K, s) and (K', s') that differ only over a small disc U such that the intersection of the tangle diagram of K with U is a contractible loop while the intersection of the tangle diagram of K' with U is empty, one has:

$$[K, s] = -(A^2 + A^{-2})[K', s'].$$

- Let $[K_1, s_1]$ and $[K_2, s_2]$ be stated skeins. Let $[K_1, s_1] = [K'_1, s'_1]$ and $[K_2, s_2] = [K'_2, s'_2]$, such that $K'_1 \subset \mathfrak{S} \times (-1, 0)$ and $K'_2 \subset \mathfrak{S} \times (0, 1)$, obtained e.g. by vertically re-scaling and translating K_1 and K_2 . Then the product $[K_1, s_1][K_2, s_2]$ of the two stated skeins in $\mathcal{S}_s^A(\Sigma, \mathcal{P})$ is defined as the stated skein $[K, s]$, where the stated V -tangle (K, s) is the union of (K'_1, s'_1) and (K'_2, s'_2) .

- A stated skein $[K, \emptyset]$ represented by a closed stated V -tangle (K, \emptyset) is said to be closed.

By forgetting the states, one can also define the (Kauffman bracket) skein algebra $\mathcal{S}^A(\Sigma, \mathcal{P})$ spanned by skeins $[K]$, which goes back to [T91]; the above is one of the several slightly different definitions of what can be called a skein algebra, suited to our purpose. It is easy to see that the subspace spanned by all closed skeins is a subalgebra, and often this subalgebra is the only focus of attention (e.g. [L17]); however, we shall see that we need the full stated skein algebra.

3.2. Square-root quantum Teichmüller space. The skein algebra $\mathcal{S}^A(\Sigma, \mathcal{P})$ is in general non-commutative, and is commutative when $A = \pm 1$. It has been observed that the family of algebras $\mathcal{S}^A(\Sigma, \mathcal{P})$, with fixed (Σ, \mathcal{P}) and A being a varying parameter, yields a version of quantization of the so-called $\mathrm{SL}_2(\mathbb{C})$ -character variety of the surface $\Sigma \setminus \mathcal{P}$ with respect to the Weil-Petersson-Goldman Poisson structure; see [BW11] for references. On the other hand, this character variety is closely related to the (enhanced) Teichmüller space equipped with the Weil-Petersson Poisson structure, whose corresponding quantum version is established by Chekhov-Fock [CF99] [F97] and by Kashaev [K98], as explained in §2.3 of the present paper. Thus it is a natural expectation that the skein algebras $\mathcal{S}^A(\Sigma, \mathcal{P})$ should be related to the quantum Teichmüller space, and a precise map relating them is what Bonahon and Wong constructed in [BW11]. However, the image of the Bonahon-Wong map lies not in the quantum Teichmüller space as presented in §2.3, but in its square-root version, which we now recall.

Following [BW11] and [H10], the Chekhov-Fock algebra \mathcal{T}_Δ^q shall be re-defined in a slightly different way than in Def.2.14. Another quantum parameter ω is introduced, and two algebras $\mathcal{T}_\Delta^\omega$ and \mathcal{T}_Δ^q will be constructed, such that \mathcal{T}_Δ^q is embedded in $\mathcal{T}_\Delta^\omega$.

Definition 3.8 ([BW11] [H10]). *Let Δ be a triangulation of a generalized marked surface (Σ, \mathcal{P}) .*

- Let t be an oriented \mathcal{P} -triangle of Δ , and let $e_{t,1}, e_{t,2}, e_{t,3}$ be the three sides of t appearing clockwise this order. In this definition, each side is viewed as a map from a closed interval into the triangle, not just as its image, hence these three sides are distinct, whether or not t is self-folded. The triangle algebra \mathcal{T}_t^ω associated to t and a nonzero complex parameter ω , is defined as the algebra over \mathbb{C} generated by three elements $\widehat{Z}_{t,1}, \widehat{Z}_{t,2}, \widehat{Z}_{t,3}$, and their inverses $\widehat{Z}_{t,1}^{-1}, \widehat{Z}_{t,2}^{-1}, \widehat{Z}_{t,3}^{-1}$, with relations

$$\widehat{Z}_{t,1}\widehat{Z}_{t,2} = \omega^2\widehat{Z}_{t,2}\widehat{Z}_{t,1}, \quad \widehat{Z}_{t,2}\widehat{Z}_{t,3} = \omega^2\widehat{Z}_{t,3}\widehat{Z}_{t,2}, \quad \widehat{Z}_{t,3}\widehat{Z}_{t,1} = \omega^2\widehat{Z}_{t,1}\widehat{Z}_{t,3},$$

together with the trivial relations $\widehat{Z}_{t,i}\widehat{Z}_{t,i}^{-1} = \widehat{Z}_{t,i}^{-1}\widehat{Z}_{t,i} = 1$, $i = 1, 2, 3$.

- The generators $\widehat{Z}_{t,1}, \widehat{Z}_{t,2}, \widehat{Z}_{t,3}$ of \mathcal{T}_t^ω are thought of as being associated to the three sides $e_{t,1}, e_{t,2}, e_{t,3}$. In particular, when the sides $e_{t,1}, e_{t,2}, e_{t,3}$ are named e, f, g , then $\widehat{Z}_{t,e}, \widehat{Z}_{t,f}, \widehat{Z}_{t,g}$ denote $\widehat{Z}_{t,1}, \widehat{Z}_{t,2}, \widehat{Z}_{t,3}$ respectively.

- Write all triangles of Δ as t_1, t_2, \dots, t_m ; so, $m = |\mathcal{F}(\Delta)|$. Consider the tensor product algebra

$$\bigotimes_{j=1}^m \mathcal{T}_{t_j}^\omega = \mathcal{T}_{t_1}^\omega \otimes \dots \otimes \mathcal{T}_{t_m}^\omega.$$

When referring to an element $Z_1 \otimes Z_2 \otimes \dots \otimes Z_m$ of $\bigotimes_{j=1}^m \mathcal{T}_{t_j}^\omega$, if some factor Z_j equals to $1 \in \mathcal{T}_{t_j}^\omega$ then we may omit that factor, provided that one can still see clearly which factor of that element lives in which factor of the tensor product algebra. The element $1 \otimes 1 \otimes \dots \otimes 1$ is denoted by 1.

- To each edge e of Δ , we associate an element \widehat{Z}_e of $\bigotimes_{j=1}^m \mathcal{T}_{t_j}^\omega$ defined as:
 - (1) If e is an internal edge that is not a self-folded edge, then e is a side of two distinct triangles t_j and t_k ; in this case, let $\widehat{Z}_e := \widehat{Z}_{t_j, e} \otimes \widehat{Z}_{t_k, e}$.
 - (2) If e is a self-folded edge of a triangle t_j , let e_1, e_2, f be the three sides of t_j appearing clockwise this order, with images of e_1 and e_2 coinciding hence forming the self-folded edge; in this case, let $\widehat{Z}_e := \omega^{-1} \widehat{Z}_{t_j, e_1} \widehat{Z}_{t_j, e_2} = \omega \widehat{Z}_{t_j, e_2} \widehat{Z}_{t_j, e_1}$.
 - (3) If e is a boundary edge, and belongs to a triangle t_j , then let $\widehat{Z}_e := \widehat{Z}_{t_j, e}$.

We also define an element \widehat{X}_e of $\bigotimes_{j=1}^m \mathcal{T}_{t_j}^\omega$ as

$$\widehat{X}_e := \widehat{Z}_e^2.$$

- Let

$$q = \omega^4.$$

Inside $\bigotimes_{j=1}^m \mathcal{T}_{t_j}^\omega$, define

$$(3.1) \quad \begin{aligned} \mathcal{T}_\Delta^\omega &:= \text{the subalgebra of } \bigotimes_{j=1}^m \mathcal{T}_{t_j}^\omega \text{ generated by } \{\widehat{Z}_e, \widehat{Z}_e^{-1} \mid e \in \Delta\}, \\ \mathcal{T}_\Delta^q &:= \text{the subalgebra of } \bigotimes_{j=1}^m \mathcal{T}_{t_j}^\omega \text{ generated by } \{\widehat{X}_e, \widehat{X}_e^{-1} \mid e \in \Delta\}, \end{aligned}$$

by a slight abuse of notation. Both are called the Chekhov-Fock algebras associated to Δ .

It is easy to observe that the generators of the Chekhov-Fock algebras satisfy the relations

$$\widehat{Z}_e \widehat{Z}_f = \omega^{2\varepsilon_{ef}} \widehat{Z}_f \widehat{Z}_e, \quad \forall e, f \in \Delta,$$

and hence

$$\widehat{X}_e \widehat{X}_f = q^{2\varepsilon_{ef}} \widehat{X}_f \widehat{X}_e, \quad \forall e, f \in \Delta.$$

Keep in mind that we have injective algebra homomorphism

$$\mathcal{T}_\Delta^q \hookrightarrow \mathcal{T}_\Delta^\omega \quad : \quad \widehat{X}_e \mapsto \widehat{Z}_e^2, \quad \forall e \in \Delta.$$

From now on, we will only use Def.3.8, instead of Def.2.14.

In order to obtain quantum coordinate change maps for the square-root generators as rational formulas, we need to consider the following.

Definition 3.9 ([BW11] [H10]). • Denote by $\widehat{Z}_{e_1}, \widehat{Z}_{e_2}, \dots, \widehat{Z}_{e_n}$ the generators of the Chekhov-Fock algebra $\mathcal{T}_\Delta^\omega$, associated to the edges of Δ labeled by e_1, e_2, \dots, e_n ; in particular, $n = |\Delta|$. A monomial in $\mathcal{T}_\Delta^\omega$ is an element $\omega^N \widehat{Z}_{e_1}^{k_1} \widehat{Z}_{e_2}^{k_2} \dots \widehat{Z}_{e_n}^{k_n}$ for some $N, k_1, k_2, \dots, k_n \in \mathbb{Z}$. A monomial $\omega^N \widehat{Z}_{e_1}^{k_1} \widehat{Z}_{e_2}^{k_2} \dots \widehat{Z}_{e_n}^{k_n}$ is said to be balanced if, for every triangle t_j of Δ , the exponents k_i of the generators \widehat{Z}_{e_i} associated to the three sides of t_j add up to an even number, where the exponent k_i for a self-folded edge e_i is counted with multiplicity 2 in this sum.

- The Chekhov-Fock (balanced) square-root algebra $\mathcal{Z}_\Delta^\omega$ is defined as the subspace of the Chekhov-Fock algebra $\mathcal{T}_\Delta^\omega$ spanned by all balanced monomials of $\mathcal{T}_\Delta^\omega$.
- The balanced square-root fraction algebra, denoted by $\widehat{\text{Frac}}(\mathcal{Z}_\Delta^\omega)$, is defined as the subalgebra of $\text{Frac}(\mathcal{T}_\Delta^\omega)$ consisting of elements that can be written as PQ^{-1} , with $P \in \mathcal{Z}_\Delta^\omega$ and $Q \in \mathcal{T}_\Delta^q \subset \mathcal{T}_\Delta^\omega$.

As mentioned in [BW11], one can check that $\widehat{\text{Frac}}(\mathcal{Z}_\Delta^\omega)$ is naturally contained in $\widehat{\text{Frac}}(\mathcal{Z}_\Delta^\omega)$. Hiatt [H10] constructed quantum coordinate change maps for the (balanced) square-root algebras extending that $\Phi_{\Delta, \Delta}^q$, for the usual Chekhov-Fock algebras, in the following sense.

Proposition 3.10 (Hiatt [H10]; see also [BW11]). *Let (Σ, \mathcal{P}) be a triangulable generalized marked surface. Let q, ω be complex numbers s.t. $q = \omega^4$. There exists an algebra isomorphism*

$$\Theta_{\Delta, \Delta'}^\omega : \widehat{\text{Frac}}(\mathcal{Z}_{\Delta'}^\omega) \rightarrow \widehat{\text{Frac}}(\mathcal{Z}_\Delta^\omega)$$

associated to each pair Δ, Δ' of triangulations of (Σ, \mathcal{P}) , satisfying:

- (1) the restriction of $\Theta_{\Delta, \Delta'}^\omega$ to $\text{Frac}(\mathcal{T}_\Delta^q)$ coincides with the map $\Phi_{\Delta, \Delta'}^q$ of Prop.2.15, and
- (2) the consistency relation holds

$$\Theta_{\Delta, \Delta''}^\omega = \Theta_{\Delta, \Delta'}^\omega \circ \Theta_{\Delta', \Delta''}^\omega$$

for each triple $\Delta, \Delta', \Delta''$ of triangulations.

Remark 3.11. *The algebra $\mathcal{Z}_\Delta^\omega$ is a quantum torus [BW17, Cor.13], and hence admits a skew-field of fractions $\text{Frac}(\mathcal{Z}_\Delta^\omega)$. It can be shown that $\text{Frac}(\mathcal{Z}_\Delta^\omega)$ coincides with $\widehat{\text{Frac}}(\mathcal{Z}_\Delta^\omega)$, simplifying the situation. Here is a sketch of proof. Viewing $\widehat{\text{Frac}}(\mathcal{Z}_\Delta^\omega)$ as a module over $\text{Frac}(\mathcal{T}_\Delta^q)$, it is a free module, as any module over a skew-field is free, and one can easily see that this module is of finite rank d . For any nonzero element x of $\widehat{\text{Frac}}(\mathcal{Z}_\Delta^\omega)$, note that $1, x, x^2, \dots, x^d$ are linearly dependent over $\text{Frac}(\mathcal{T}_\Delta^q)$, hence $\sum_{i=1}^d c_i x^i = 0$ for some $c_0, \dots, c_d \in \text{Frac}(\mathcal{T}_\Delta^q)$ that are not all zero. Since x is not a zero divisor, c_0 must be nonzero, hence can be assumed to be 1. It follows $1 = -(c_1 + c_2 x + \dots + c_d x^{d-1})x$, showing that x is invertible. So $\widehat{\text{Frac}}(\mathcal{Z}_\Delta^\omega)$ is a skew-field, hence coincides with the skew-field of fractions of $\mathcal{Z}_\Delta^\omega$.*

As pointed out in [BW11], this map $\Theta_{\Delta, \Delta'}^\omega$ is much better understood in terms of operators on Hilbert spaces. This operator-theoretic viewpoint will be made clear and explicit e.g. in an upcoming work [KS] (see [S20b] for an attempt).

Similarly for the case of $\text{Frac}(\mathcal{T}_\Delta^q)$, by the square-root quantum Teichmüller space we mean the quotient of the disjoint union of all $\widehat{\text{Frac}}(\mathcal{Z}_\Delta^\omega)$ by the equivalence relation given by the identifications $\Theta_{\Delta, \Delta'}^\omega$. Also, as an analog of Def.2.16, the ring of square-root quantum regular functions can be defined as

$$(3.2) \quad \mathbf{L}_\Delta^\omega := \bigcap_{\Delta'} \Theta_{\Delta, \Delta'}^\omega(\mathcal{Z}_{\Delta'}^\omega) \subset \mathcal{Z}_\Delta^\omega \subset \widehat{\text{Frac}}(\mathcal{Z}_\Delta^\omega) \subset \text{Frac}(\mathcal{T}_\Delta^\omega),$$

by a slight abuse of notation with the previously defined symbol \mathbf{L}_Δ^q . As $\Theta_{\Delta, \Delta'}^\omega$ induces a natural isomorphism from $\mathbf{L}_{\Delta'}^\omega$ to \mathbf{L}_Δ^ω , we may denote \mathbf{L}_Δ^ω for all Δ 's collectively as $\mathbf{L}^\omega = \mathbf{L}_{(\Sigma, \mathcal{P})}^\omega$, or understand this situation as $\mathbf{L}^\omega = \bigsqcup_{\Delta} \mathbf{L}_\Delta^\omega / \sim$, where \sim is the equivalence relation coming from the maps $\Theta_{\Delta, \Delta'}^\omega$.

3.3. Quantum trace map. If we restrict our attention only to triangulable generalized marked surfaces (Σ, \mathcal{P}) with empty boundary, i.e. punctured surfaces, we are ready to state the result of Bonahon-Wong, namely a map from the skein algebra of this surface to the square-root quantum Teichmüller space. In the meantime, one of the major defining properties of this map is the cutting/gluing property, which is a certain compatibility that holds when cutting the surface (together with a skein) along a \mathcal{P} -arc, i.e. an edge of some triangulation. In order to fully reflect this property, it is not just a luxury but rather a must, that we state Bonahon-Wong's result in complete generality for any triangulable generalized marked surface, instead of only for punctured surfaces without boundary.

Bonahon and Wong [BW11] described the process of gluing surfaces and skeins along two boundary arcs of a not-necessarily-connected surface. To conveniently attain uniqueness of such a process, we instead formulate everything in terms of cutting.

Definition and Lemma 3.12 (cutting construction). *Let (Σ, \mathcal{P}) be a generalized marked surface, not necessarily connected nor triangulable. Let b be a \mathcal{P} -arc in Σ , and assume that the interior \mathring{b} of b lies in the interior $\Sigma \setminus \partial\Sigma$ of the surface.*

- *Let (Σ', \mathcal{P}') be the unique (up to diffeomorphism) generalized marked surface obtained from (Σ, \mathcal{P}) by cutting along b . In particular, there is a natural (gluing) map $\Sigma' \rightarrow \Sigma$, whose restriction to \mathcal{P}' yields a correspondence $\mathcal{P}' \rightarrow \mathcal{P}$. The pre-image of b under $\Sigma' \rightarrow \Sigma$ is the union of two boundary arcs of (Σ', \mathcal{P}') , denoted by b_1 and b_2 . The gluing map $\Sigma' \rightarrow \Sigma$ restricts to a diffeomorphism $\Sigma' \setminus (b_1 \cup b_2) \rightarrow \Sigma \setminus b$.*

- *Suppose that K is a tangle in $(\Sigma \setminus \mathcal{P}) \times (-1, 1)$ satisfying the following conditions:*

- (C1) *K is transverse to $b \times (-1, 1)$, and*
- (C2) *the framing of K at each point in $K \cap (b \times (-1, 1))$ is upward vertical.*

The above process of cutting along b uniquely yields a tangle K' in $(\Sigma' \setminus \mathcal{P}') \times (-1, 1)$, equipped with a map $K' \rightarrow K$. The number of pre-images of $x \in K$ under $K' \rightarrow K$ is 2 if $x \in b$, and is 1 otherwise.

- Suppose further that (Σ, \mathcal{P}) is triangulable, and that Δ is a triangulation of (Σ, \mathcal{P}) such that b is an internal edge of Δ , i.e. $b \in \mathring{\Delta}$. The process of cutting along b uniquely yields a triangulation Δ' of (Σ', \mathcal{P}') .

- In each of the above three situations, we say that the new object (for (Σ', \mathcal{P}')) is obtained from the former by cutting along b .

Lemma 3.13 ([BW11]). *Suppose the situation of Def.3.12, and $\omega \in \mathbb{C}^*$. The triangles of Δ are naturally in one-to-one correspondence with those of Δ' , and the sides of each triangle of Δ are naturally in one-to-one correspondence with those of the corresponding triangle of Δ' . Hence we have a natural isomorphism $\bigotimes_{t \in \mathcal{F}(\Delta)} \mathcal{T}_t^\omega \rightarrow \bigotimes_{t' \in \mathcal{F}(\Delta')} \mathcal{T}_{t'}^\omega$ of algebras, and this restricts to an injective algebra homomorphism*

$$(3.3) \quad \mathcal{Z}_\Delta^\omega \rightarrow \mathcal{Z}_{\Delta'}^\omega$$

between the Chekhov-Fock (balanced) square-root algebras.

Definition 3.14 ([BW11]). *Let (Σ, \mathcal{P}) be a generalized marked surface, not necessarily triangulable. Let $A \in \mathbb{C}^*$, and let $\mathcal{S}_s^A(\Sigma, \mathcal{P})$ be the stated skein algebra defined in Def.3.7.*

- Suppose that (Σ', \mathcal{P}') is obtained from (Σ, \mathcal{P}) by cutting along a \mathcal{P} -arc b of Σ , and that a tangle K' in $(\Sigma' \setminus \mathcal{P}') \times (-1, 1)$ is obtained from a tangle K in $(\Sigma \setminus \mathcal{P}) \times (-1, 1)$ through this cutting process, as described in Def.3.12. We say that the states $s' : \partial K' \rightarrow \{+, -\}$ and $s : \partial K \rightarrow \{+, -\}$ for these tangles are compatible if the following holds: for each $x' \in \partial K'$, denote by $x \in K$ the image of x' under the map $K' \rightarrow K$ in Def.3.12

(CS1) If $x \in \partial K$, then $s'(x') = s(x)$;

(CS2) If $x \notin \partial K$, i.e. $x \in b$, so that the pre-image of x under $K' \rightarrow K$ is $\{x', x''\}$, then $s'(x') = s'(x'')$.

We can finally state the result of Bonahon and Wong in full generality.

Proposition 3.15 (the quantum trace map; the main theorem of [BW11]). *Let A and ω be nonzero complex numbers s.t.*

$$A = \omega^{-2}.$$

Then there is a unique family of algebra homomorphisms

$$(3.4) \quad \text{Tr}_\Delta^\omega = \text{Tr}_{(\Sigma, \mathcal{P}); \Delta}^\omega : \mathcal{S}_s^A(\Sigma, \mathcal{P}) \rightarrow \mathcal{Z}_\Delta^\omega$$

from the stated skein algebra to the Chekhov-Fock (balanced) square-root algebra, defined for each triangulable generalized marked surface (Σ, \mathcal{P}) and each triangulation Δ of (Σ, \mathcal{P}) , satisfying:

- (1) (Cutting Property¹) Suppose that (Σ', \mathcal{P}') , Δ' and K' are related to (Σ, \mathcal{P}) , Δ and K as in Def.3.12, i.e. the former are obtained by cutting the latter along an internal edge of Δ . Then for each state s for the tangle K , we have

$$\text{Tr}_{(\Sigma, \mathcal{P}); \Delta}^\omega([K, s]) = \sum_{\text{compatible } s'} \text{Tr}_{(\Sigma', \mathcal{P}'); \Delta'}^\omega([K', s']),$$

where the sum is over all states s' for the tangle K' for the cut surface (Σ', \mathcal{P}') that are compatible with s in the sense of Def.3.14. The left-hand-side, which is a priori an element of $\mathcal{Z}_\Delta^\omega$, is viewed as an element of $\mathcal{Z}_{\Delta'}^\omega$ via the embedding map in eq.(3.3).

- (2) (Elementary Cases) Suppose that (Σ, \mathcal{P}) is a non-self-folded triangle, i.e. Σ is diffeomorphic to a closed disc, and \mathcal{P} consists of three marked points on the boundary; let Δ denote its unique triangulation. If $[K, s] \in \mathcal{S}_s^A(\Sigma, \mathcal{P})$ is a stated skein where the boundary-ordered tangle diagram of K is one of the two cases (a) and (b) in Fig.3, we have:

¹Bonahon and Wong [BW11] call this ‘State Sum Property’




FIGURE 3. Boundary-ordered tangle diagrams for elementary cases of a skein in a triangle (arrows on the sides indicate the boundary-orientation (Def.2.1))

- (a) Suppose that \widehat{Z}_1 and \widehat{Z}_2 are the generators of the Chekhov-Fock algebra $\mathcal{T}_\Delta^\omega$ of eq.(3.1) associated to the edges of Δ containing the endpoints x_1 and x_2 in Fig.3(a), respectively. In particular, $\widehat{Z}_1\widehat{Z}_2 = \omega^2\widehat{Z}_2\widehat{Z}_1$. Then,

$$(3.5) \quad \text{Tr}_{(\Sigma, \mathcal{P}); \Delta}^\omega([K, s]) = \begin{cases} 0 & \text{if } s(x_1) = - \text{ and } s(x_2) = +, \\ \left[\widehat{Z}_1^{s(x_1)} \widehat{Z}_2^{s(x_2)} \right]_{\text{Weyl}} & \text{otherwise,} \end{cases}$$

where a sign $\varepsilon = \pm$ in the exponent means ± 1 respectively.

- (b) For K as in Fig.3(b), one has

$$\text{Tr}_{(\Sigma, \mathcal{P}); \Delta}^\omega([K, s]) = \begin{cases} 0 & \text{if } s(x_1) = s(x_2), \\ -\omega^{-5} & \text{if } s(x_1) = + \text{ and } s(x_2) = -, \\ \omega^{-1} & \text{if } s(x_1) = - \text{ and } s(x_2) = +. \end{cases}$$

For a triangulated generalized marked surface, cutting along all the internal edges of the triangulation yields disjoint union of non-self-folded triangles. Also, using the skein relations repeatedly, any stated skein $[K, s] \in \mathcal{S}_s^A(\Sigma, \mathcal{P})$ can be written as a linear combination of stated skeins whose tangle diagrams have no crossing at all. Along this line, one can show that the properties of $\text{Tr}_{(\Sigma, \mathcal{P}); \Delta}^\omega$ in Prop.3.15 completely determine the values of the maps $\text{Tr}_{(\Sigma, \mathcal{P}); \Delta}^\omega$, if the existence of these maps is assumed.

A crucial property of Bonahon-Wong quantum trace is its compatibility with the quantum coordinate-change maps.

Proposition 3.16 ([BW11]). *Let Δ, Δ' be triangulations of a triangulable generalized marked surface (Σ, \mathcal{P}) . Then we have*

$$\text{Tr}_\Delta^\omega = \Theta_{\Delta, \Delta'}^\omega \circ \text{Tr}_{\Delta'}^\omega,$$

where $\Theta_{\Delta, \Delta'}^\omega : \widehat{\text{Frac}}(\mathcal{Z}_{\Delta'}^\omega) \rightarrow \widehat{\text{Frac}}(\mathcal{Z}_\Delta^\omega)$ is the (square-root) quantum coordinate-change isomorphism in Prop.3.10. More precisely, for each stated skein $[K, s] \in \mathcal{S}_s^A(\Sigma, \mathcal{P})$, with $A = \omega^{-2}$, the balanced Laurent polynomial $\text{Tr}_{\Delta'}^\omega([K, s]) \in \mathcal{Z}_{\Delta'}^\omega$ is sent by $\Theta_{\Delta, \Delta'}^\omega$ to the balanced Laurent polynomial $\text{Tr}_\Delta^\omega([K, s]) \in \mathcal{Z}_\Delta^\omega$.

In particular, the images of Bonahon-Wong quantum trace map are balanced Laurent polynomials in the square-root generators for *any* triangulation Δ , hence belong to \mathbf{L}_Δ^ω defined in eq.(3.2).

We now briefly introduce Allegretti-Kim's idea [AK17] of using the Bonahon-Wong quantum trace to obtain a solution to the quantum ordering problem discussed in §2.4. Consider a \mathcal{P} -knot γ in a triangulable generalized marked surface (Σ, \mathcal{P}) ; that is, γ is a simple loop in the surface $\mathfrak{S} = \Sigma \setminus \mathcal{P}$. Suppose that γ is not a contractible loop. Then γ represents a nontrivial element $[\gamma]$ of $\pi_1(\mathfrak{S})$, say, if we choose a basepoint. Given any point ρ of the Teichmüller space $\mathcal{T}(\Sigma, \mathcal{P})$ defined by eq.(2.1), one considers the monodromy $\rho([\gamma])$, which is an element of $\text{PSL}(2, \mathbb{R})$ defined up to conjugation in $\text{PSL}(2, \mathbb{R})$; then $|\text{trace}(\rho([\gamma]))|$, the absolute value of the trace of this monodromy, is a well-defined real number. This provides a smooth function $\mathbb{I}(\gamma)$ on $\mathcal{X}^+(\Sigma, \mathcal{P})$, whose value at each point (ρ, O) is defined to be $|\text{trace}(\rho([\gamma]))|$. It turns out that, for *any* chosen ideal triangulation Δ of (Σ, \mathcal{P}) , the function $\mathbb{I}(\gamma)$ is a (positive-)integer-coefficient Laurent polynomial in the square-roots $X_e^{1/2}$ of the exponentiated shear coordinate functions of edges e of Δ . Based on these functions which naturally arise geometrically, Fock and Goncharov [FG06] proposed a basis of the ring of all universally Laurent functions, i.e. functions that are Laurent polynomials in the (square-roots of the) exponentiated shear coordinate functions for every ideal triangulation, where this basis is enumerated by '(even) integral laminations' on the surface

\mathfrak{S} , which are multicurves with integer weights subject to certain condition. For a proof that this basis indeed spans all universally Laurent functions, see [FG06] [S20a] [GHK15]. One can view a simple loop γ as a special example of an integral lamination. In order to construct a quantum version $\widehat{\mathbb{I}}^\omega(\gamma)$ that deforms the classical function $\mathbb{I}(\gamma)$, we first lift the curve γ living in the surface \mathfrak{S} to a tangle in the 3d manifold $\mathfrak{S} \times (-1, 1)$, the thickening of \mathfrak{S} .

Definition 3.17. *Let (Σ, \mathcal{P}) be a generalized marked surface. Let γ be a simple closed curve in $\mathfrak{S} = \Sigma \setminus \mathcal{P}$, i.e. a closed \mathcal{P} -link in Σ .*

Denote by K_γ^c the tangle in $\mathfrak{S} \times (-1, 1)$ obtained as the lift of γ at a constant elevation $c \in (-1, 1)$ with upward vertical framing everywhere. We call K_γ^c a constant-elevation lift of γ . We may denote K_γ^c by K_γ without specifying c .

Note that K_γ is well-defined up to isotopy, for a given γ . An easy observation:

Lemma 3.18. *A closed tangle K in $\mathfrak{S} \times (-1, 1)$ is isotopic to a tangle whose tangle diagram in Σ has no crossing at all if and only if K is isotopic to a constant-elevation lift K_γ of a simple closed curve γ in $\Sigma \setminus \mathcal{P}$. ■*

Note that K_γ is a closed tangle, hence a closed V-tangle. Choose any ideal triangulation Δ of (Σ, \mathcal{P}) , and apply the Bonahon-Wong quantum trace map in eq.(3.4) to the stated skein $[K_\gamma, \emptyset] \in \mathcal{S}_s^A(\Sigma, \mathcal{P})$; the resulting element of $\mathcal{Z}_\Delta^\omega$ is the Allegretti-Kim quantum element associated to $\mathbb{I}(\gamma)$:

$$(3.6) \quad \widehat{\mathbb{I}}_\Delta^\omega(\gamma) := \text{Tr}_{(\Sigma, \mathcal{P}); \Delta}^\omega([K_\gamma, \emptyset]) \in \mathbf{L}_\Delta^\omega \subset \mathcal{Z}_\Delta^\omega.$$

Treatment for more general integral laminations need several more crucial ideas. See [AK17] for this, and also for various favorable properties enjoyed by these quantum elements; see also [CKKO20] for an important positivity property. To present a couple of examples of its nice properties, we have the quantum mutation compatibility $\widehat{\mathbb{I}}_\Delta^\omega(\gamma) = \Theta_{\Delta, \Delta'}^\omega(\widehat{\mathbb{I}}_{\Delta'}^\omega(\gamma))$, and it is relatively easy to see that $\widehat{\mathbb{I}}_\Delta^\omega(\gamma)$ indeed recovers the classical function $\mathbb{I}(\gamma)$ when $\omega = 1$ (see e.g. [BW11]). In particular, the assignment $\mathbb{I}(\gamma) \mapsto \widehat{\mathbb{I}}^\omega(\gamma) = \widehat{\mathbb{I}}_\Delta^\omega(\gamma) = \text{Tr}_{(\Sigma, \mathcal{P}); \Delta}^\omega([K_\gamma, \emptyset])$ provides a partial answer to the quantum ordering problem mentioned in §2.4; see [AK17] for a full answer, which requires algebraic manipulations including Chebyshev polynomials, as well as certain control of parity of powers of monomials.

3.4. Biangle quantum trace. Although Propositions 3.15 and 3.16 completely describe and determine the Bonahon-Wong quantum trace map, they are not very convenient when it comes to actual computation of the values. For any given stated skein $[K, s] \in \mathcal{S}_s^A(\Sigma, \mathcal{P})$, there is a more direct algorithm that enables us to compute the quantum trace $\text{Tr}_\Delta^\omega([K, s])$ called the ‘state-sum formula’, which we shall recall in §3.5. As a preliminary step for that formula, we first recall the quantum trace for biangles, in the present subsection.

Recall that a *biangle* B is a generalized marked surface (Σ, \mathcal{P}) , with Σ diffeomorphic to a closed disc, which in particular has one boundary component, where \mathcal{P} consists of two marked points on the boundary. As noted in Def.2.4, it is not triangulable, hence there is no Bonahon-Wong quantum trace map that Prop.3.15 associates to it; in particular, the quantum Teichmüller space is not defined. However, its stated skein algebra makes sense, because Definition 3.7 applies. Bonahon and Wong [BW11] defined and studied a quantum trace map for biangles, separately from Prop.3.15.

Recall that in our notations, we have $\partial B = \partial \Sigma \setminus \mathcal{P}$, and ∂B is a disjoint union of two boundary arcs of B . Let’s say that a \mathcal{P} -arc in B is an *internal arc* if its interior is contained in the interior of Σ . Recall that each generalized marked surface (Σ, \mathcal{P}) is equipped with an orientation on Σ . Hence, by a biangle we automatically mean an oriented biangle. Notice that any two biangles are diffeomorphic.

For an (oriented) biangle B viewed as a generalized marked surface as above, choose an internal arc b connecting the two marked points of B . Then, cutting B along b yields a unique (up to diffeomorphism) generalized marked surface (Σ', \mathcal{P}') as described in Def.3.12. One easily observes that (Σ', \mathcal{P}') is a disjoint union of two biangles. Here is an analog of Prop.3.15 for biangles:

Proposition 3.19 (quantum trace for biangles; [BW11]). *Let $A, \omega \in \mathbb{C}^*$ satisfy $A = \omega^{-2}$. Then there is a unique family of algebra homomorphisms*

$$\text{Tr}_B^\omega : \mathcal{S}_s^A(B) \rightarrow \mathbb{C}$$

defined for all (oriented) biangles B , such that

- (1) (*Cutting Property*) Let $B = (\Sigma, \mathcal{P})$ be a biangle, and $[K, s] \in \mathcal{S}_s^A(B)$ be a stated skein for B . Let (Σ', \mathcal{P}') be the generalized marked surface obtained by cutting B along an internal arc of B connecting the two marked points, as described in Def.3.12. Then (Σ', \mathcal{P}') is disjoint union of two oriented biangles $B_1 = (\Sigma_1, \mathcal{P}_1)$ and $B_2 = (\Sigma_2, \mathcal{P}_2)$, and the tangle $K \subseteq (\Sigma \setminus \mathcal{P}) \times (-1, 1)$ yields tangles $K_1 \subseteq (\Sigma_1 \setminus \mathcal{P}_1) \times (-1, 1)$ and $K_2 \subseteq (\Sigma_2 \setminus \mathcal{P}_2) \times (-1, 1)$ by this cutting process. Then one has

$$\mathrm{Tr}_B^\omega([K, s]) = \sum_{\text{compatible } s_1, s_2} \mathrm{Tr}_{B_1}^\omega([K_1, s_1]) \mathrm{Tr}_{B_2}^\omega([K_2, s_2]),$$

where the sum is over all pairs of states $s_1 : \partial K_1 \rightarrow \{+, -\}$ and $s_2 : \partial K_2 \rightarrow \{+, -\}$ that comprise states $s' : \partial K' \rightarrow \{+, -\}$ of (Σ', \mathcal{P}') that are compatible with $s : \partial K \rightarrow \{+, -\}$ in the sense of Def.3.14.

- (2) (*Elementary Cases*) For a single biangle B , if $[K, s] \in \mathcal{S}_s^A(B)$ is a stated skein for B consisting of one component, and the boundary-ordered tangle diagram of K is one of the two cases (I) and (II) in Fig.4:

(I) One has

$$\mathrm{Tr}_B^\omega([K, s]) = \begin{cases} 1 & \text{if } s(x) = s(y), \\ 0 & \text{if } s(x) \neq s(y); \end{cases}$$

(II) One has

$$\mathrm{Tr}_B^\omega([K, s]) = \begin{cases} 0 & \text{if } s(y_1) = s(y_2), \\ -\omega^{-5} & \text{if } s(y_1) = - \text{ and } s(y_2) = +, \\ \omega^{-1} & \text{if } s(y_1) = + \text{ and } s(y_2) = -. \end{cases}$$




FIGURE 4. Some boundary-ordered tangle diagrams in a biangle (arrows on the boundary arcs indicate the boundary-orientation (Def.2.1))

For any given stated skein for a single biangle B , by using skein relations one can resolve it to a linear combination of elementary stated skeins dealt with in Prop.3.19(2) and therefore the value under Tr_B^ω can be computed. One caveat is that even when the stated tangle diagram of a stated skein $[K, s]$ over B is without crossing, the value of the biangle quantum trace can still be complicated, instead of being just products of the above elementary cases (I) and (II). For later use, we list some examples:

Lemma 3.20 (Biangle quantum trace on remaining elementary tangles). *Let A, ω and B be as in Prop.3.19. For a stated skein $[K, s] \in \mathcal{S}_s^A(B)$ falling into one of the following cases, we have:*

- (1) *If the boundary-ordered tangle diagram of K is as in case (III) of Fig.4, one has ([BW11, Lem.14])*

$$(3.7) \quad \mathrm{Tr}_B^\omega([K, s]) = \begin{cases} 0 & \text{if } s(x_1) = s(x_2), \\ \omega & \text{if } s(x_1) = - \text{ and } s(x_2) = +, \\ -\omega^5 & \text{if } s(x_1) = + \text{ and } s(x_2) = -. \end{cases}$$

- (2) *In case (IV) of Fig.4 when the tangle diagram consists of two disjoint parallel lines, if we let*

$$(3.8) \quad \epsilon = \begin{cases} +1 & \text{if the vertical orderings are such that } x_1 \succ x_2 \text{ and } y_2 \succ y_1, \\ -1 & \text{if } x_2 \succ x_1 \text{ and } y_1 \succ y_2, \end{cases}$$

and $s(x_1) = \varepsilon_1$, $s(x_2) = \varepsilon_2$, $s(y_1) = \varepsilon'_1$ and $s(y_2) = \varepsilon'_2$, then one has:

$$(3.9) \quad \mathrm{Tr}_B^\omega([K, s]) = \begin{cases} \omega^{2\epsilon} & \text{if } \varepsilon_1 = \varepsilon_2 = \varepsilon'_1 = \varepsilon'_2, \\ \omega^{2\epsilon} - \omega^{-6\epsilon} & \text{if } \varepsilon_1 = \varepsilon'_2 = + \text{ and } \varepsilon_2 = \varepsilon'_1 = -, \\ \omega^{-2\epsilon} & \text{if } \varepsilon_1 = \varepsilon'_1 \neq \varepsilon'_2 = \varepsilon_2, \\ 0 & \text{otherwise.} \end{cases}$$

- (3) In case of Fig.4(V), the value of $\mathrm{Tr}_B^\omega([K, s])$ is given by eq.(3.9) with $\epsilon = -1$ and $s(x_1) = \varepsilon_2$, $s(x_2) = \varepsilon_1$, $s(y_1) = \varepsilon'_1$ and $s(y_2) = \varepsilon'_2$.
- (4) In case of Fig.4(VI), the value of $\mathrm{Tr}_B^\omega([K, s])$ is given by eq.(3.9) with $\epsilon = +1$ and $s(x_1) = \varepsilon_1$, $s(x_2) = \varepsilon_2$, $s(y_1) = \varepsilon'_2$ and $s(y_2) = \varepsilon'_1$.

Genuinely simple basic cases can be conveniently described if one adapts the peculiar picture convention of Bonahon-Wong [BW11, §3.5] for stated boundary-ordered diagrams in a surface \mathfrak{S} , which stipulates that the vertical ordering on each boundary arc b of \mathfrak{S} should match the horizontal ordering with respect to a chosen orientation on b . If the stated boundary-ordered tangle diagram in a biangle B consists of disjoint parallel lines under this Bonahon-Wong picture convention where one boundary arc is given the boundary-orientation and the other the opposite of the boundary-orientation, then the value $\mathrm{Tr}_B^\omega([K, s])$ is either 0 or 1. Since we do not use the Bonahon-Wong convention, we translate this observation as follows:

Lemma 3.21 (biangle factor of a skein whose diagram consists of (strongly) parallel lines). *Let $[K, s]$ be a stated skein over a biangle $B = (\Sigma, \mathcal{P})$. Let $\omega \in \mathbb{C}^*$ and $A = \omega^{-2}$. Let e and e' be the two boundary arcs of B , constituting the boundary of B . Suppose that the V-tangle $K \subset (\Sigma \setminus \mathcal{P}) \times (-1, 1)$ satisfies:*

- (1) *the stated (boundary-ordered) tangle diagram of K in B consists of disjoint parallel lines (under our picture convention, not necessarily under Bonahon-Wong convention);*
- (2) *the ordering on the segments of K induced by the elevations of the points of $\partial K \cap (e \times (0, 1))$ (i.e. the endpoints of the segments of K over e) coincides with that induced by the elevations of the points of $\partial K \cap (e' \times (0, 1))$.*

If, for every segment of K , its two endpoints are assigned the same sign by s , then $\mathrm{Tr}_B^\omega([K, s]) = 1$. Otherwise, $\mathrm{Tr}_B^\omega([K, s]) = 0$.

Proof. Let k_1, \dots, k_r be the segments of K , i.e. the components of K . For each $j = 1, \dots, r$, let $I_j \subset (-1, 1)$ be the image of k_j under the projection $(\Sigma \setminus \mathcal{P}) \times (-1, 1) \rightarrow (-1, 1)$ to the second factor, i.e. the collection of elevations of the points of k_j . By condition (2), we see that K can be V-isotoped so that I_1, \dots, I_r are mutually disjoint. So one can rename these segments so that the sequence k_1, \dots, k_r is arranged in the increasing order of elevations. Since $\mathrm{Tr}_B^\omega : \mathcal{S}_s^A(B) \rightarrow \mathbb{C}$ is an algebra homomorphism (Prop.3.19), where the product in $\mathcal{S}_s^A(B)$ is given by the superposition operation, it follows

$$\mathrm{Tr}_B^\omega([K, s]) = \mathrm{Tr}_B^\omega([k_1, s_1]) \mathrm{Tr}_B^\omega([k_2, s_2]) \cdots \mathrm{Tr}_B^\omega([k_r, s_r]).$$

Each $\mathrm{Tr}_B^\omega([k_j, s_j])$ falls into the case of Prop.3.19.(2)(a), hence the claim follows. \blacksquare

3.5. State-sum formula. The sought-for state-sum formula for $\mathrm{Tr}_\Delta^\omega$ requires the notion of ‘split’ ideal triangulation $\widehat{\Delta}$ of an ideal triangulation Δ , where each edge of Δ is replaced by a biangle. Then the complexity of a skein caused by elevations shall be pushed to biangles by isotopy.

Definition 3.22 ([BW11]). *Let Δ be a triangulation of a generalized marked surface (Σ, \mathcal{P}) , and let $\mathfrak{S} = \Sigma \setminus \mathcal{P}$. Recall that Δ is a collection of \mathcal{P} -arcs in Σ satisfying certain conditions. Denote the edges of Δ by e_1, e_2, \dots, e_n , and the triangles of Δ by t_1, t_2, \dots, t_m .*

• For each edge e_i of Δ choose a \mathcal{P} -arc e'_i in Σ that is \mathcal{P} -isotopic to e_i , so that no two members of the collection

$$\widehat{\Delta} := \Delta \cup \{e'_1, \dots, e'_n\}$$

intersect in $\Sigma \setminus \mathcal{P}$. Call this $\widehat{\Delta}$ a split \mathcal{P} -triangulation (or, split ideal triangulation, or split triangulation) associated to the triangulation Δ . For each i , the region bounded by e_i and e'_i is called a biangle B_i . The triangles formed by $\widehat{\Delta}$ are in one-to-one correspondence with the triangles of Δ , and we denote them by $\widehat{t}_1, \dots, \widehat{t}_m$, correspondingly.

• A V-tangle K in $\mathfrak{S} \times (-1, 1)$ is said to be in a good position (with respect to $\widehat{\Delta}$) if it satisfies (P1) of §3.1 and all of the following:

- (GP1) For each constituent edge e of $\widehat{\Delta}$, K is transverse to $e \times (-1, 1)$. In particular, $K \cap (e \times (-1, 1))$ has at most finitely many elements;
- (GP2) For every triangle \widehat{t}_j of $\widehat{\Delta}$, $K \cap (\widehat{t}_j \times (-1, 1))$ consists of finitely many disjoint arcs, each of which is contained in a constant elevation surface $\widehat{t}_j \times *$ and joins two distinct components of $\partial\widehat{t}_j \times (-1, 1)$, where each component of $\partial\widehat{t}_j$ is a side of \widehat{t}_j minus the vertices of $\partial\widehat{t}_j$;
- (GP3) For every triangle \widehat{t}_j of $\widehat{\Delta}$, the components of $K \cap (\widehat{t}_j \times (-1, 1))$ lie at mutually distinct elevations;
- (GP4) None of the crossings of K lie over an edge of $\widehat{\Delta}$.

Note that every triangle of $\widehat{\Delta}$ has three distinct sides, even if the corresponding triangle of Δ is self-folded. In particular, in (GP2) above, for each triangle \widehat{t}_j of $\widehat{\Delta}$, the number of components of $\partial\widehat{t}_j$, hence also the number of components of $\partial\widehat{t}_j \times (-1, 1)$, is always three. We added (GP4) for convenience.

Lemma 3.23 (from [BW11]). *A VH-tangle K in $\mathfrak{S} \times (-1, 1)$ can be VH-isotoped to a VH-tangle in a good position.*

For a V-tangle in a good position, the elevation change occurs only over the biangles. Note that a V-tangle being in a good position does *not* guarantee that its tangle diagram on a triangle of $\widehat{\Delta}$ has no crossing.

We find it convenient to define some more words, both for stating Bonahon-Wong's construction and for later sections of the present paper.

Definition 3.24. *Let (Σ, \mathcal{P}) , \mathfrak{S} , Δ , $\widehat{\Delta}$, e_i , e'_i , t_j and \widehat{t}_j be as in Def.3.22; in particular, i runs through $1, \dots, n$ and j runs through $1, \dots, m$. Let K be a V-tangle in $\mathfrak{S} \times (-1, 1)$ in a good position with respect to $\widehat{\Delta}$. Let $E_{\widehat{\Delta}} := \bigcup_{i=1}^n (e_i \cup e'_i)$. Then K is divided by $E_{\widehat{\Delta}} \times (-1, 1)$ into (tangle) segments. Projection on Σ of each segment of K is also called a (tangle) segment. The boundary points of a segment are called $\widehat{\Delta}$ -junctures of K (or just junctures of K), or endpoints of that segment.*

A $\widehat{\Delta}$ -juncture-state of K is a map $J : \{\widehat{\Delta}\text{-junctures of } K\} \rightarrow \{+, -\}$ assigning a sign to each juncture.

The word 'junctures' could have been defined just as elements of $K \cap (E_{\widehat{\Delta}} \times (-1, 1))$, or their projections in Σ . The points of ∂K and their projections are also examples of junctures.

Proposition 3.25 (state-sum formula of Bonahon-Wong quantum trace; [BW11, §6]). *Let (Σ, \mathcal{P}) , \mathfrak{S} , Δ , $\widehat{\Delta}$, e_i , e'_i , B_i , t_j and \widehat{t}_j be as in Def.3.22, and K be a V-tangle in $\mathfrak{S} \times (-1, 1)$ in a good position with respect to $\widehat{\Delta}$. Let $s : \partial K \rightarrow \{+, -\}$ be a state for the tangle K . Let $A, \omega \in \mathbb{C}^*$ satisfy $A = \omega^{-2}$.*

Let $J : \{\widehat{\Delta}\text{-junctures of } K\} \rightarrow \{+, -\}$ be a $\widehat{\Delta}$ -juncture-state of K . For each triangle \widehat{t}_j of $\widehat{\Delta}$, let $K_j := K \cap (\widehat{t}_j \times (-1, 1))$. Then $[K_j, J|_{\partial K_j}]$ is a stated skein for the triangle \widehat{t}_j , where $J|_{\partial K_j}$ is the restriction of J to ∂K_j . Let $k_{j,1}, \dots, k_{j,l_j}$ be the components of K_j , i.e. the tangle segments of K over the triangle \widehat{t}_j , in order of increasing elevation. Then each $[k_{j,\alpha}, J|_{\partial k_{j,\alpha}}]$ is a stated skein for the triangle \widehat{t}_j , where $J|_{\partial k_{j,\alpha}}$ is the restriction of J to $\partial k_{j,\alpha}$. Then the element $\text{Tr}_{\widehat{t}_j}^\omega([k_{j,\alpha}, J|_{\partial k_{j,\alpha}}])$ of the triangle algebra $\mathcal{T}_{\widehat{t}_j}^\omega$ is defined via Prop.3.15(2)(a), and the element $\text{Tr}_{\widehat{t}_j}^\omega([K_j, J|_{\partial K_j}])$ of $\mathcal{T}_{\widehat{t}_j}^\omega$ is given by

$$(3.10) \quad \text{Tr}_{\widehat{t}_j}^\omega([K_j, J|_{\partial K_j}]) = \text{Tr}_{\widehat{t}_j}^\omega([k_{j,1}, J|_{\partial k_{j,1}}]) \text{Tr}_{\widehat{t}_j}^\omega([k_{j,2}, J|_{\partial k_{j,2}}]) \cdots \text{Tr}_{\widehat{t}_j}^\omega([k_{j,l_j}, J|_{\partial k_{j,l_j}}]) \in \mathcal{T}_{\widehat{t}_j}^\omega.$$

Via the natural map $\mathcal{T}_{\widehat{t}_j}^\omega \rightarrow \mathcal{T}_{t_j}^\omega$ induced by the correspondence of the sides of the triangles, this element $\text{Tr}_{\widehat{t}_j}^\omega([K_j, J|_{\partial K_j}])$ can be viewed as an element of $\mathcal{T}_{t_j}^\omega$.

For each biangle $B_i = (\Sigma_i, \mathcal{P}_i)$ which is bounded by e_i and e'_i , let $L_i := K \cap ((\Sigma_i \setminus \mathcal{P}_i) \times (-1, 1))$. Then $[L_i, J|_{\partial L_i}]$ is a stated skein for the biangle B_i , where B_i is viewed as a generalized marked surface of its own, and $J|_{\partial L_i}$ is the restriction of J to ∂L_i . Define the Bonahon-Wong term for the $\widehat{\Delta}$ -juncture-state J of the V-tangle K as

$$(3.11) \quad \text{BW}_{\widehat{\Delta}}^\omega(K; J) := \left(\prod_{i=1}^n \text{Tr}_{B_i}^\omega([L_i, J|_{\partial L_i}]) \right) \left(\bigotimes_{j=1}^m \text{Tr}_{\widehat{t}_j}^\omega([K_j, J|_{\partial K_j}]) \right) \in \bigotimes_{j=1}^m \mathcal{T}_{t_j}^\omega,$$

where the numbers $\text{Tr}_{B_i}^\omega([L_i, J|_{\partial L_i}]) \in \mathbb{C}$ are given by Prop.3.19. Then $\text{BW}_{\widehat{\Delta}}^\omega(K; J) \in \mathcal{Z}_{\widehat{\Delta}}^\omega \subset \mathcal{T}_{\widehat{\Delta}}^\omega$.

Finally, one has

$$\mathrm{Tr}_\Delta^\omega([K, s]) = \sum_{J: J|_{\partial K} = s} \mathrm{BW}_\Delta^\omega(K; J) \in \mathcal{Z}_\Delta^\omega \subset \mathcal{T}_\Delta^\omega \subset \bigotimes_{j=1}^m \mathcal{T}_{t_j}^\omega,$$

where the sum is over all $\widehat{\Delta}$ -junction-states J that restricts to the given state s at ∂K .

The above proposition is what one can practically use for actual computation of the values of the Bonahon-Wong quantum trace.

4. GABELLA'S QUANTUM HOLONOMY

4.1. Branched double cover surface. Quantization of the trace-of-monodromy functions (for closed curves) on the Teichmüller space, namely the quantum ordering problem mentioned in §2.4, is also of interest to physicists, as the coefficients of the monomials of the quantum version of the trace-of-monodromy correspond to the so-called ‘framed protected spin characters’ in physics. A very interesting solution to this quantum ordering problem is given by Gabella [G17]. Gabella’s construction of quantum holonomy is based on the works of Gaiotto-Moore-Neitzke [GMN13] [GMN14] and Galakhov-Longhi-Moore [GLM15]. Part of the main ideas of these works are the processes called ‘non-abelianization’ and ‘abelianization’, which relate a GL_N -bundle of a surface Σ to an abelian bundle of a certain N -fold branched (i.e. ramified) covering of Σ . In our case, $N = 2$. We assume Σ is connected. We start with the description of a branched double cover of the surface Σ , and some basic constructions about it.

Recall that a *branched double cover* $\pi : \widetilde{\Sigma} \rightarrow \Sigma$, with a finite branching set $\mathcal{V} \subset \Sigma$, is a continuous map such that the restriction $\pi' : \widetilde{\Sigma} \setminus \pi^{-1}(\mathcal{V}) \rightarrow \Sigma \setminus \mathcal{V}$ of π is an ordinary double covering and the restriction of π onto $\pi^{-1}(\mathcal{V})$ is bijective. Given the branching set \mathcal{V} , branched double covers π of Σ are in one-to-one correspondence with ordinary double covers π' of $\Sigma \setminus \mathcal{V}$, which are classified by subgroups of index 2 of the fundamental group of $\Sigma \setminus \mathcal{V}$, or cohomology classes in $H^1(\Sigma \setminus \mathcal{V}; \mathbb{Z}/2\mathbb{Z})$. Here we will identify $\mathbb{Z}/2\mathbb{Z}$ with the multiplicative group $\{+, -\} = \{+1, -1\}$.

Definition 4.1 (branched double cover). *Let (Σ, \mathcal{P}) be a triangulable generalized marked surface and Δ a triangulation of (Σ, \mathcal{P}) . For each triangle t of Δ , choose a point v_t in the interior of t . These points are called the branch points. Denote by \mathcal{V} the set of all branch points. The manifold $\Sigma \setminus \mathcal{V}$ deformation retracts to the CW-complex consisting of vertices \mathcal{P} and edges of the triangulation Δ . The 1-cocycle assigning to every edge of this complex the element $-$ of the group $\{+, -\}$ defines a 1-cohomology class in $H^1(\Sigma \setminus \mathcal{V}; \mathbb{Z}/2\mathbb{Z})$, and hence defines a branched double cover*

$$\pi : \widetilde{\Sigma}_\Delta \rightarrow \Sigma$$

with the branching set \mathcal{V} .

Remark 4.2. *The above is a special case of a more general theory of branched N -fold cover of Σ , which corresponds to a ‘Seiberg-Witten curve’ in physics [G12] [GMN12]. See these references, as well as [GMN13] [GMN14] [G17], for descriptions using branch cuts.*

A basic observation is that every continuous path $\alpha : [0, 1] \rightarrow \Sigma \setminus \mathcal{V}$ has two *continuous lifts* $[0, 1] \rightarrow \widetilde{\Sigma}_\Delta \setminus \pi^{-1}(\mathcal{V})$ in the branched double cover. Lifting paths in Σ to paths in $\widetilde{\Sigma}_\Delta$ is a crucial part of the non-abelianization process. To keep track of the lifted paths, we need the following.

Definition and Lemma 4.3 (states of lifts of marked points). *Let (Σ, \mathcal{P}) , Δ and $\pi : \widetilde{\Sigma}_\Delta \rightarrow \Sigma$ be as in Def.4.1. Let $\widetilde{\mathcal{P}} = \pi^{-1}(\mathcal{P})$ be the lifts of the marked points. Define a state function $s = s_\Delta : \widetilde{\mathcal{P}} \rightarrow \{+, -\}$ as follows. First choose a point in $\widetilde{\mathcal{P}}$ and assign value $+$ for the function s at this point. There is a unique extension $s : \widetilde{\mathcal{P}} \rightarrow \{+, -\}$ such that if $p, p' \in \widetilde{\mathcal{P}}$ are connected by a continuous lift of an edge in Δ , then they have different s value. We will say that $p \in \widetilde{\mathcal{P}}$ is on the sheet $s(p) \in \{+, -\}$.*

The state $s : \widetilde{\mathcal{P}} \rightarrow \{+, -\}$ depends on the choice of Δ , and there are two possible choices of states s , for a given Δ ; just choose one s . We now consider lifting paths in Σ to paths in $\widetilde{\Sigma}_\Delta$. A crucial example of the branched double cover is the case when the base surface (Σ, \mathcal{P}) is a triangle.

Definition 4.4. *Let t be a non-self-folded triangle, viewed as a generalized marked surface (Σ, \mathcal{P}) ; that is, Σ is diffeomorphic to a closed disc and \mathcal{P} consists of three points on $\partial\Sigma$. Let Δ be the unique ideal triangulation of t , and denote $\widetilde{\Sigma}_\Delta$ by \widetilde{t} . Let x_0 and x_1 be points of $\partial t = \partial\Sigma \setminus \mathcal{P}$ lying in distinct sides*




FIGURE 5. Gabella lifts in a triangle, and their states

of t , say b_0 and b_1 respectively. A Gabella lift of the ordered pair (x_0, x_1) is any proper embedding $\tilde{\gamma} : [0, 1] \rightarrow \tilde{t} \setminus \pi^{-1}(\{v_t\})$ such that $\pi(\tilde{\gamma}(i)) = x_i$ for each $i = 0, 1$, or its image together with the orientation.

For a given pair (x_0, x_1) , there are two distinct paths in $t \setminus \{v_t\}$ from x to y , up to isotopy in $t \setminus \{v_t\}$. Hence, up to isotopy in \tilde{t} , there are four distinct Gabella lifts of the pair (x_0, x_1) , which can be conveniently parametrized by the signs at points x_0 and x_1 as follows.

Definition 4.5. Let $(\Sigma, \mathcal{P}) = t$ and \tilde{t} be as in Def.4.4. Assume $\tilde{\gamma} : [0, 1] \rightarrow \tilde{t} \setminus \pi^{-1}(\{v_t\})$ is a Gabella lift of the pair (x_0, x_1) of points of ∂t lying in distinct sides b_0 and b_1 of t . For each $i = 0, 1$, among the two continuous lifts in \tilde{t} of b_i , let \tilde{b}_i be the one that contains $\tilde{\gamma}(i)$. Let $\tilde{\gamma}'(i)$ be the velocity vector of $\tilde{\gamma}$ at $\tilde{\gamma}(i)$. At the point $\tilde{\gamma}(i) \in \tilde{b}_i$ choose a tangent vector to \tilde{b}_i that forms with $\tilde{\gamma}'(i)$ the positive orientation of the surface, and travel along \tilde{b}_i in the direction of this tangent vector until one reaches a point \tilde{p} of $\tilde{\mathcal{P}}$. Define the state of $\tilde{\gamma}$ at x_i to be $s(x_i) := s(\tilde{p})$. The resulting map $s = s_{\tilde{\gamma}} : \{x_0, x_1\} \rightarrow \{+, -\}$ is the state of the Gabella lift $\tilde{\gamma}$ of the pair (x_0, x_1) .

See Fig.5 for examples. It is easy to observe the following, from the definition.

Lemma 4.6. Given any map $s : \{x_0, x_1\} \rightarrow \{+, -\}$, there is a unique, up to isotopy in \tilde{t} , Gabella lift of the pair (x_0, x_1) whose state coincides with s . ■

A Gabella lift $\tilde{\gamma}$ of (x_0, x_1) can be thought of as a lift of a path $\gamma = \pi \circ \tilde{\gamma}$ in t from x_0 to x_1 . Meanwhile, we should eventually be dealing with the thickenings of t and \tilde{t} . For our purposes, it suffices to consider the following:

Definition 4.7. Let $(\Sigma, \mathcal{P}) = t$ and \tilde{t} be as in Def.4.4.

- A horizontal triangle segment over t is an oriented VH -tangle $k : [0, 1] \rightarrow t \times (-1, 1)$ in the thickening of t living at a constant elevation $c \in (-1, 1)$ (i.e. image of k lies in $t \times \{c\}$), equipped with the upward vertical framing, such that $\partial k \neq \emptyset$ and that the elements of ∂k lie in distinct components of $\partial t \times (-1, 1)$, as in (GP2) of Def.3.22; in particular, $k(u) = (\gamma(u), c)$ for some path $\gamma : [0, 1] \rightarrow t$.

- By a Gabella lift of a horizontal triangle segment k we mean any path $\tilde{k} : [0, 1] \rightarrow \tilde{t} \times (-1, 1)$ of the form $\tilde{k}(u) = (\tilde{\gamma}(u), c)$, where $\tilde{\gamma}$ is a Gabella lift of the pair $(\gamma(0), \gamma(1))$.

- The state of the Gabella lift \tilde{k} of k is $s = s_{\tilde{k}} : \{k(0), k(1)\} \rightarrow \{+, -\}$ induced by $s_{\tilde{\gamma}}$ of Def.4.5.

- Let \tilde{k} be a Gabella lift of k . Let e_1, e_2, e_3 be the sides of t , appearing clockwise in ∂t in this order. Suppose the endpoints of γ lies in e_i and e_{i+1} (with $e_4 = e_1$). If the value of the state $s_{\tilde{\gamma}}$ at the endpoint of γ in e_i is $-$ and that at the endpoint in e_{i+1} is $+$, then \tilde{k} is said to be non-admissible. Otherwise, \tilde{k} is admissible.

In Fig.5, the only non-admissible Gabella lift is the leftmost curve in the picture on the right. Given a horizontal triangle segment k over t , there are four Gabella lifts of k up to isotopy in $t \times (-1, 1)$, parametrized by the four possible states $s : \{k(0), k(1)\} \rightarrow \{+, -\}$ at the endpoints of k . Exactly one of these four Gabella lifts is non-admissible.

Now we deal with the case of any triangulable generalized marked surface (Σ, \mathcal{P}) .

Definition 4.8. *Let (Σ, \mathcal{P}) be a triangulable generalized marked surface, and Δ an ideal triangulation of (Σ, \mathcal{P}) . Let $\mathfrak{S} = \Sigma \setminus \mathcal{P}$. Let $\widehat{\Delta}$ be a split ideal triangulation of Δ (Def.3.22). Let K be an oriented VH-tangle in $\mathfrak{S} \times (-1, 1)$ in a good position with respect to $\widehat{\Delta}$ (Def.3.22), so that the tangle segments of K (Def.3.24) over each triangle of $\widehat{\Delta}$ is a horizontal tangle segment; call these triangle segments of K .*

A Gabella lift \widetilde{K} (in $\widetilde{\Sigma}_\Delta \times (-1, 1)$) of K is the choice of a Gabella lift of each triangle segment of K . We say that \widetilde{K} is admissible if the chosen Gabella lift of each triangle segment of K is admissible. An isotopy of Gabella lifts of K is a simultaneous isotopy of the Gabella lifts of the triangle segments.

We now briefly describe physicists' ideas of abelianization and non-abelianization. The original object of our study is the enhanced Teichmüller space $\mathcal{X}^+(\Sigma, \mathcal{P})$ of a triangulable generalized marked surface (Σ, \mathcal{P}) , defined in Def.2.11. One standard way of studying this space is to view it as the moduli space of PSL(2, \mathbb{R})-local systems on $\mathfrak{S} = \Sigma \setminus \mathcal{P}$ satisfying some conditions [FG06], together with certain data at the 'asymptotic boundary points' \mathcal{P} . Recall that, for a Lie group G , a G -local system on a manifold means a principal G -bundle on the manifold together with a flat G -connection on it. The abelianization and non-abelianization processes of Gaiotto-Moore-Neitzke [GMN13] build a correspondence between the moduli space of (certain) GL_N -local systems on the original surface \mathfrak{S} and the moduli space of (certain) GL_1 -local systems on a branched N -fold cover $\widetilde{\mathfrak{S}}$ of \mathfrak{S} . The GL_N parallel transport map of a path γ in \mathfrak{S} corresponds to the sum of (formal) GL_1 parallel transports of lifts $\widetilde{\gamma}$ of this path in the branched cover $\widetilde{\mathfrak{S}}$. Not all lifts contribute to this sum, but only the ones satisfying an admissibility condition do, which corresponds to our admissibility condition when $N = 2$. The term associated to $\widetilde{\gamma}$ is written as a product of GL_1 parallel transports of some elementary paths in $\widetilde{\mathfrak{S}}$, hence can be viewed as a monomial in some variables. Upon quantization, these monomials should be enhanced to non-commutative monomials, and an appropriate quantum ordering must be chosen for the resulting (Laurent) polynomial to satisfy favorable properties.

One answer was suggested by Galakhov, Longhi and Moore [GLM15]; namely, consider the Weyl-ordered non-commutative monomial times a power of q to the net sum of signs of self-intersections of the lifted path $\widetilde{\gamma}$. In view of the way how they determined the signs, this net sum of signs is really the writhe of a framed path in the thickened 3-manifold $\widetilde{\mathfrak{S}} \times (-1, 1)$ given the 'always going up' elevations that projects to $\widetilde{\gamma}$ in $\widetilde{\mathfrak{S}}$. In particular, a path γ is first lifted to a framed path in the 3d manifold $\mathfrak{S} \times (-1, 1)$ with always-going-up elevations, and then the quantum GL_n parallel transport is constructed using writhes of lifted paths in $\widetilde{\mathfrak{S}} \times (-1, 1)$. So, this method was insufficient to deal with the quantum parallel transport along a *closed* path γ , because it can't be lifted to a closed framed path in $\mathfrak{S} \times (-1, 1)$ with always-going-up elevations. Once we choose a starting point of γ , we lift it to an always-going-up path in $\mathfrak{S} \times (-1, 1)$, and then need to add a 'going down' path in the end in order to get a closed path. Gabella's contribution [G17] is the consideration of a certain complex-number correction factor associated to this closing-up part, in order to make the final result to be independent of the choice of starting basepoint. This correction factor, which Gabella refers to as an *R-matrix*, is heavily inspired by the biangle factor of Bonahon-Wong [BW11] which was reviewed in §3.4 of the present paper.

In fact, we enhance Gabella's construction and describe it for any *general* oriented VH-tangle K in the thickened surface $\mathfrak{S} \times (-1, 1)$, not just for a (simple) loop in \mathfrak{S} nor a certain special closed tangle in $\mathfrak{S} \times (-1, 1)$ that has always-going-up elevations except at a small part. For now, assume that K is in a good position with respect to some split ideal triangulation $\widehat{\Delta}$; later, our main result will allow us to drop this assumption. Gabella's quantum holonomy along such K in a good position will be given as a sum of certain terms, over all admissible Gabella lifts \widetilde{K} of K up to isotopy. One might want to understand each \widetilde{K} as a framed tangle in $\widetilde{\Sigma}_\Delta \times (-1, 1)$ that lifts K , but in general, \widetilde{K} cannot be a continuous path in $\widetilde{\Sigma}_\Delta \times (-1, 1)$ and must have discontinuities over biangles of $\widehat{\Delta}$. However, we still refer to \widetilde{K} as a Gabella lift of K in $\widetilde{\Sigma}_\Delta \times (-1, 1)$, for convenience.

In pictures, a Gabella lift \widetilde{K} is drawn in \mathfrak{S} as follows. In each biangle B of $\widehat{\Delta}$, just draw the boundary-ordered tangle diagram for $K \cap (B \times (-1, 1))$. In each triangle \widehat{t} of $\widehat{\Delta}$, isotope \widetilde{K} so that the image of \widetilde{K} under the projection $\pi \times \text{id} : \widetilde{\Sigma}_\Delta \times (-1, 1) \rightarrow \mathfrak{S} \times (-1, 1)$ satisfies (P1) and (P2) of §3.1 over \widehat{t} , hence yielding a boundary-ordered tangle diagram in \widehat{t} . We also indicate the values of the states at

the endpoints of each triangle segment over \hat{t} . These data forms a *diagram of \tilde{K}* . In particular, it is a boundary-ordered tangle diagram in \mathfrak{S} together with some extra data. By a *crossing of \tilde{K}* we mean a self-intersection of the image of \tilde{K} under the projection $\tilde{\Sigma}_\Delta \times (-1, 1) \rightarrow \tilde{\Sigma}_\Delta$, and we indicate this in the diagram of \tilde{K} drawn in \mathfrak{S} as before, using broken lines. A self-intersection of the diagram of \tilde{K} that is not a crossing of \tilde{K} is drawn without broken lines; see Fig.10.

4.2. Biangle quantum holonomy via operator invariant. The strategy of Gabella [G17] to construct a quantum holonomy is similar to Bonahon-Wong in the following sense: first define the quantum holonomy for a biangle, and then define the quantum holonomy for a triangulated surface via a state-sum formula. We first focus on biangles. We note that, over a biangle, Gabella [G17, §5.3–§5.4] describes the values of the (biangle) quantum holonomy only for simple examples of oriented VH-tangles for which he refers to as ‘R-matrix’ and ‘cup/cap’. Gabella mentions that his R-matrix associated to an oriented tangle in a biangle consisting of a single crossing can be interpreted as the R-matrix appearing in the representation theory of the quantum group $\mathcal{U}_q(\mathfrak{gl}_N)$, where we can put $N = 2$ for now. This strongly hints that the biangle quantum holonomy of Gabella is related to the Reshetikhin-Turaev invariant of tangles [RT90] [T16], associated to a representation of a quantum group. We will eventually confirm this expectation later in the present paper.

We first review and settle some basic setting for constructing operator invariants of tangles in a biangle, by mostly following the contents of Ohtsuki’s book [O02]. Translating into our language requires us to consider a biangle with fixed labeling of boundary arcs. First, recall from Def.2.4 and §3.4 that a *biangle B* is an example of a generalized marked surface (Σ, \mathcal{P}) . In particular, it is an oriented surface diffeomorphic to a closed disc with two marked points on the boundary, and its boundary $\partial B = \partial\Sigma \setminus \mathcal{P}$ is a disjoint union of two boundary arcs. We will need to work with a following version of biangle.

Definition 4.9 (directed biangle). • A *direction* of a biangle B is a bijective map

$$(4.1) \quad \text{dir} = \text{dir}_B : \{\text{the two boundary arcs of } B\} \rightarrow \{\text{in, out}\}.$$

The boundary arc mapping to in is called the *inward* boundary arc b_{in} , and the one mapping to out the *outward* boundary arc b_{out} .

- A *directed biangle \vec{B}* is a pair (B, dir) consisting of a biangle B and a direction dir of B .
- In a directed biangle $\vec{B} = (B, \text{dir})$, the outward boundary arc b_{out} is given the boundary-orientation (Def.2.1) on b_{out} coming from the surface orientation on B . The inward boundary arc b_{in} is given the orientation opposite to the boundary-orientation on b_{in} . These orientations are depicted in Fig.7.

Any two directed biangles can be identified by a diffeomorphism preserving the direction. Note that once a direction is chosen on a biangle, there is one of the two marked points of B that both the oriented boundary arcs b_{in} and b_{out} point toward. So the choice of a direction is equivalent to the choice of such a distinguished marked point of B . To get Ohtsuki’s pictures, one should rotate our pictures by 90 degrees clockwise.

Definition 4.10. • Let $\mathcal{D}(\vec{B})$ be the set of all equivalence classes $[D]$ of boundary-ordered oriented tangle diagrams D (Def.3.3) in \vec{B} , where two boundary-ordered oriented tangle diagrams are defined to be *equivalent* if one can be obtained from the other by a sequence of isotopies of boundary-ordered oriented tangle diagrams and moves (M1) in §3.1, i.e. the framed Reidemeister moves I, II and III.

- Let $\mathcal{D}_s(\vec{B})$ be the set of all equivalence classes $[D, s]$ of stated boundary-ordered oriented tangle diagrams (D, s) , where the equivalence is defined analogously.

By Prop.3.5(4) we see that $\mathcal{D}(\vec{B})$ is in one-to-one correspondence with the set of all VH-isotopy classes of VH-tangles in the thickening of a directed biangle. As any two directed biangles \vec{B} and \vec{B}' are identified by a diffeomorphism preserving the directions, the sets $\mathcal{D}(\vec{B})$ and $\mathcal{D}(\vec{B}')$ can be naturally identified. We now study some basic operations on $\mathcal{D}(\vec{B})$.

Definition 4.11. Let \vec{B} be a directed biangle, with the underlying biangle denoted by $B = (\Sigma, \mathcal{P})$. Identify $\mathfrak{S} = \Sigma \setminus \mathcal{P}$ with $[0, 1] \times \mathbb{R}$ by choosing a diffeomorphism such that b_{in} mapsto $\{1\} \times \mathbb{R}$ and b_{out} to $\{0\} \times \mathbb{R}$, where the orientations of b_{in} and b_{out} match the usual increasing orientation of \mathbb{R} . The \mathbb{R} -coordinate is called a *horizontal* coordinate.




FIGURE 6. Tensor product and composition of tangle diagrams in a directed biangle

We say that a collection D_1, \dots, D_n of tangle diagrams in \vec{B} are horizontally-disjoint if the images of them under the second projection $[0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ are mutually disjoint. We say D_i is horizontally higher than D_j if the \mathbb{R} -coordinates of points of D_i are bigger than those of D_j .

We say that a collection D_1, \dots, D_n of boundary-ordered tangle diagrams in \vec{B} are vertically-disjoint if for each pair i, j of distinct indices in $\{1, \dots, n\}$, either D_i is vertically higher than D_j , i.e. on each of the two boundary arcs b any element of $\partial_b D_i$ has higher vertical ordering than any element of $\partial_b D_j$, or D_j is vertically higher than D_i .

A collection of boundary-ordered tangle diagrams in \vec{B} is said to be completely-disjoint if it is horizontally-disjoint and vertically-disjoint.

For a tangle diagram D in a directed biangle \vec{B} , we write

$$(4.2) \quad \partial_{\text{in}} D = \partial_{b_{\text{in}}} D \quad \text{and} \quad \partial_{\text{out}} D = \partial_{b_{\text{out}}} D,$$

where b_{in} and b_{out} are inward and outward boundary arcs of \vec{B} respectively.

Definition 4.12 (tensor product and composition of oriented tangle diagrams in a directed biangle). Let \vec{B} be a directed biangle, and let $[D_1], [D_2] \in \mathcal{D}(\vec{B})$.

- Let D'_1 and D'_2 be boundary-ordered oriented tangle diagrams in \vec{B} that are isotopic to D_1 and D_2 respectively, such that D'_1 and D'_2 are completely-disjoint and D'_1 is horizontally higher than D'_2 . Let D be the union of D'_1 and D'_2 , with D'_1 being set to be vertically higher than D'_2 ; see Fig.6. Define the tensor product of the equivalence classes of boundary-ordered oriented tangle diagrams $[D_1]$ and $[D_2]$ in \vec{B} as $[D_1] \otimes [D_2] := [D]$.

- Suppose that $\partial_{\text{in}} D_1$ and $\partial_{\text{out}} D_2$ have the same cardinality, that the unique bijection $\partial_{\text{in}} D_1 \rightarrow \partial_{\text{out}} D_2$ that preserves the horizontal orderings also preserves the vertical orderings, and that this bijection is compatible with the orientations on D_1 and D_2 in the sense that it sends sinks to sources and vice versa; we then say that $[D_1]$ is composable with $[D_2]$. Now, say that D_1 is a boundary-ordered oriented tangle diagram in a directed biangle \vec{B}_1 , and D_2 is one in \vec{B}_2 . Let \vec{B}' be the directed biangle obtained by gluing \vec{B}_1 and \vec{B}_2 along the inward boundary arc of \vec{B}_1 and the outward boundary arc of \vec{B}_2 respecting their orientations, so that the gluing map restricts to the above bijection $\partial_{\text{in}} D_1 \rightarrow \partial_{\text{out}} D_2$. Let D be the boundary-ordered oriented tangle diagram in \vec{B}' obtained as the union of the images of D_1 and D_2 under the gluing map; see Fig.6. Define the composition of the equivalence classes of boundary-ordered oriented tangle diagrams $[D_1]$ and $[D_2]$ in a directed biangle as $[D_1] \circ [D_2] := [D]$.

It is well known that any oriented tangle diagram in a directed biangle \vec{B} can be obtained from certain elementary ones by applying a sequence of tensor products and compositions; we formulate the result in terms of boundary-ordered oriented tangle diagrams.

Definition 4.13 (elementary tangle diagrams in a directed biangle; e.g. [O02]). • A boundary-ordered oriented tangle diagram D in a directed biangle \vec{B} is said to be elementary if it is one of the 10 cases in Fig.7. We say the corresponding equivalence class $[D]$ is elementary.

Lemma 4.14. Any $[D] \in \mathcal{D}(\vec{B})$ can be obtained from elementary classes in $\mathcal{D}(\vec{B})$ by applying a sequence of tensor products and compositions.




FIGURE 7. Elementary boundary-ordered oriented tangle diagrams in a directed biangle (orientations on boundary arcs b_{out} and b_{in} are as in Def.4.9)

We are now ready to formulate Gabella's biangle quantum holonomy, as matrix elements of an operator invariant. From now on let

$$(4.3) \quad V := \text{the 2-dimensional } \mathbb{C} \text{ vector space, equipped with the ordered basis } \{\xi_+, \xi_-\}.$$

We note that, in Gabella's notation [G17], the basis vectors ξ_+ and ξ_- correspond to the symbols s_1 and s_2 . For $D \in \mathcal{D}(\vec{B})$ we will consider $V^{\otimes |\partial_{\text{in}} D|}$ and $V^{\otimes |\partial_{\text{out}} D|}$, where the tensor factors are thought to be ordered according to the (increasing) horizontal orderings on $\partial_{\text{in}} D$ and $\partial_{\text{out}} D$; for example, in case of Fig.4(IV), we have $V^{\otimes |\partial_{\text{in}} D|} = V \otimes V$, with the first factor corresponding to the endpoint x_1 and the second factor to x_2 . We set $V^{\otimes 0} := \mathbb{C}$.

Proposition 4.15 (Gabella's biangle quantum holonomy packaged as an operator invariant). *There exists a unique map G assigning to each $[D] \in \mathcal{D}(\vec{B})$ a linear operator*

$$G([D]) : V^{\otimes |\partial_{\text{in}} D|} \rightarrow V^{\otimes |\partial_{\text{out}} D|}$$

that satisfies

$$G([D_1] \circ [D_2]) = G([D_1]) \circ G([D_2]), \quad G([D_1] \otimes [D_2]) = G([D_1]) \otimes G([D_2]),$$

and whose values at elementary classes $[D] \in \mathcal{D}(\vec{B})$ are:

- (GB1) (identity) if D is as in Fig.7(I) or (VI), then $G([D]) = \text{id} : V \rightarrow V$;
- (GB2) (cup) if D is as in Fig.7(II) or (VII), then $G([D]) : \mathbb{C} \rightarrow V \otimes V$ sends $1 \in \mathbb{C}$ to $\xi_+ \otimes \xi_- - \omega^4 \xi_- \otimes \xi_+$;
- (GB3) (cap) if D is as in Fig.7(III) or (VIII), then $G([D]) : V \otimes V \rightarrow \mathbb{C}$ sends $\xi_+ \otimes \xi_-$ to $-\omega^{-4}$ and $\xi_- \otimes \xi_+$ to 1 while other basis vectors to zero;
- (GB4) (height exchange) if D is as in Fig.7(IV), then the map $G([D]) : V \otimes V \rightarrow V \otimes V$ is given on the basis vectors as:

$$(4.4) \quad G([D]) : \xi_i \otimes \xi_j \mapsto \begin{cases} \xi_+ \otimes \xi_- + (\omega^4 - \omega^{-4}) \xi_- \otimes \xi_+, & \text{if } i = + \text{ and } j = -, \\ \xi_i \otimes \xi_j, & \text{otherwise,} \end{cases}$$

while if D is as in Fig.7(XI), then the map $G([D]) : V \otimes V \rightarrow V \otimes V$ is given as the inverse of the map in eq.(4.4);

- (GB5) (positive crossing) if D is as in Fig.7(V), then $G([D]) : V \otimes V \rightarrow V \otimes V$ is given on the basis vectors as:

$$(4.5) \quad G([D]) : \xi_i \otimes \xi_j \mapsto \begin{cases} \omega^{-4} \xi_i \otimes \xi_j, & \text{if } i = j, \\ \xi_- \otimes \xi_+, & \text{if } i = + \text{ and } j = -, \\ \xi_+ \otimes \xi_- + (\omega^{-4} - \omega^4) \xi_- \otimes \xi_+, & \text{if } i = - \text{ and } j = +; \end{cases}$$

(GB6) (negative crossing) if D is as in Fig.7(X), then $G([D]) : V \otimes V \rightarrow V \otimes V$ is given as the inverse of the map in eq.(4.5).

What will be actually used in the construction of Gabella's quantum holonomy over the entire triangulated surface are matrix elements.

Definition 4.16 (Basis of $V^{\otimes M}$ and bilinear form). For each ordered sequence $\vec{\varepsilon} = \{\varepsilon_1, \dots, \varepsilon_M\}$ in $\{+, -\}$, define the basis vector of $V^{\otimes M}$ as

$$\xi^{\vec{\varepsilon}} := \xi_{\varepsilon_1} \otimes \xi_{\varepsilon_2} \otimes \dots \otimes \xi_{\varepsilon_M} \in V \otimes V \otimes \dots \otimes V = V^{\otimes M}.$$

Define a symmetric bilinear form $\langle \cdot, \cdot \rangle$ on $V^{\otimes M}$ as $\langle \xi^{\vec{\varepsilon}}, \xi^{\vec{\varepsilon}'} \rangle = \delta_{\vec{\varepsilon}, \vec{\varepsilon}'}$, the Kronecker delta.

Definition 4.17 (Gabella's biangle quantum holonomy as matrix elements of G). Let \vec{B} be a directed biangle, with the underlying biangle being $B = (\Sigma, \mathcal{P})$, with $\mathfrak{S} = \Sigma \setminus \mathcal{P}$. Let $\omega \in \mathbb{C}^*$. Let $[K, s]$ be the VH-isotopy class of a stated oriented VH-tangle in $\mathfrak{S} \times (-1, 1)$.

Denote by $[D, s] \in \mathcal{D}_s(\vec{B})$ the stated boundary-ordered tangle diagram of $[K, s]$. Let $\partial_{\text{in}}(D) = \{x_1, x_2, \dots, x_{|\partial_{\text{in}} D|}\}$ and $\partial_{\text{out}}(D) = \{y_1, \dots, y_{|\partial_{\text{out}} D|}\}$, arranged in the respective (increasing) horizontal orderings. Define the inward/outward basis vector for s as

$$(4.6) \quad \xi_{\text{in}}^s := \xi_{s(x_1)} \otimes \xi_{s(x_2)} \otimes \dots \otimes \xi_{s(x_{|\partial_{\text{in}} D|})} \in V^{\otimes |\partial_{\text{in}} D|}, \quad \xi_{\text{out}}^s := \xi_{s(y_1)} \otimes \dots \otimes \xi_{s(y_{|\partial_{\text{out}} D|})} \in V^{\otimes |\partial_{\text{out}} D|},$$

where in case $\partial_{\text{in}} D = \emptyset$ or $\partial_{\text{out}} D = \emptyset$, we let $\xi_{\text{in}}^s = 1$ or $\xi_{\text{out}}^s = 1$ respectively. Let $G([D, s]) \in \mathbb{C}$ be the matrix element of the linear map $G([D]) : V^{\otimes |\partial_{\text{in}} D|} \rightarrow V^{\otimes |\partial_{\text{out}} D|}$ of Prop.4.15 for these basis vectors, i.e.

$$G([D, s]) := \langle \xi_{\text{out}}^s, G([D]) \xi_{\text{in}}^s \rangle$$

Define the Gabella biangle quantum holonomy as the map $\text{TrHol}_{\vec{B}}^{\omega}$ assigning to each $[K, s]$ the value:

$$\text{TrHol}_{\vec{B}}^{\omega}([K, s]) := G([D, s]) \in \mathbb{C}.$$

For example, if D is as in Prop.4.15(GB2) with $y_1 \succ y_2$, $s(y_1) = -$ and $s(y_2) = +$, then $\text{TrHol}_{\vec{B}}^{\omega}([K, s]) = -\omega^4$, while if D is as in Prop.4.15(GB4) with $s(x_1) = +$, $s(x_2) = -$, $s(y_1) = -$ and $s(y_2) = +$, then $\text{TrHol}_{\vec{B}}^{\omega}([K, s]) = \omega^4 - \omega^{-4}$.

The following useful observations can be checked in a straightforward manner; we will provide a proof of it at the end of §5.5, using the relationship with the Bonahon-Wong biangle quantum trace.

Lemma 4.18. *The Gabella biangle quantum holonomy satisfies:*

- (1) (direction independence) $\text{TrHol}_{\vec{B}}^{\omega}([K, s])$ is independent of the choice of the direction on B , so we can define

$$\text{TrHol}_B^{\omega}([K, s]) := \text{TrHol}_{\vec{B}}^{\omega}([K, s]).$$

- (2) (charge conservation) A state s of a tangle diagram D in \vec{B} or of a VH-tangle in the thickening of \vec{B} is said to be charge-preserving iff the net sum of signs assigned by s to elements of $\partial_{\text{in}} D$ equals that for $\partial_{\text{out}} D$ (where \pm are viewed as ± 1). We have $\text{TrHol}_B^{\omega}([K, s]) = 0$ for any state s that is not charge-preserving.

Above is a rigorous treatment via operator invariants of what Gabella meant in [G17]. In particular, one can verify by inspection that the values of TrHol_B^{ω} at the elementary stated oriented VH-tangles defined as above coincide with Gabella's values in [G17, §5] except for the cups and caps. The values at cups and caps originally assigned by Gabella in [G17, eq.(5.7)] should be corrected to our values given by Def.4.17 and Prop.4.15(GB2)–(GB3), in order to guarantee the well-definedness (i.e. isotopy invariance) of Gabella's quantum holonomy for the entire surface which will be constructed in the next subsection. We will fully justify our construction of TrHol_B^{ω} , e.g. prove Prop.4.15, later.

4.3. Quantum holonomy. We now describe Gabella's quantum holonomy [G17] associated to an oriented VH-tangle K in $\mathfrak{S} \times (-1, 1)$. We first define a Gabella lift of a stated oriented VH-tangle (K, s) , and enumerate their isotopy classes by the collection of states for the Gabella lifts of triangle segments, which we package as a $\widehat{\Delta}$ -juncture-state of K .




FIGURE 8. Sign of a crossing

Definition 4.19. Let (Σ, \mathcal{P}) , \mathfrak{S} , Δ and $\widehat{\Delta}$ be as in Def.4.8. Let K be an oriented VH-tangle in $\mathfrak{S} \times (-1, 1)$ in a good position with respect to $\widehat{\Delta}$. Let \widetilde{K} be a Gabella lift in $\widetilde{\Sigma}_\Delta \times (-1, 1)$ of K . The $\widehat{\Delta}$ -juncture-state $J_{\widetilde{K}}$ of K whose restriction to the endpoints of each triangle segment k of K coincides with the state $s_{\widetilde{k}}$ of the Gabella lift \widetilde{k} of k determined by \widetilde{K} is the $\widehat{\Delta}$ -juncture-state of K associated to Gabella lift \widetilde{K} .

If s is a state of K , by a Gabella lift of the stated oriented VH-tangle (K, s) we mean a Gabella lift \widetilde{K} of K whose associated $\widehat{\Delta}$ -juncture-state $J_{\widetilde{K}}$ restricts to s at ∂K .

Lemma 4.20. Let (K, s) be a stated oriented VH-tangle in $\mathfrak{S} \times (-1, 1)$ in a good position with respect to $\widehat{\Delta}$. The correspondence $\widetilde{K} \mapsto J_{\widetilde{K}}$ yields a bijection between the set of all isotopy classes of Gabella lifts of K and the set of all $\widehat{\Delta}$ -juncture-states of K . Gabella lifts of (K, s) correspond to $\widehat{\Delta}$ -juncture-states of K restricting to s at ∂K .

Definition 4.21. For a $\widehat{\Delta}$ -juncture-state J of K , denote by \widetilde{K}^J a Gabella lift of K whose associated $\widehat{\Delta}$ -juncture-state coincides with J .

A $\widehat{\Delta}$ -juncture-state J of K is said to be admissible if \widetilde{K}^J is an admissible Gabella lift of K (Def.4.8), that is, if none of the triangle segments of \widetilde{K}^J fall into the case of Fig.3(a) with $J(x_1) = -$ and $J(x_2) = +$.

Remark 4.22. Notice that the state-sum formula of the Bonahon-Wong quantum trace, as seen in §3.5, was written as a sum over all $\widehat{\Delta}$ -juncture-states of K . One may observe that the terms corresponding to non-admissible $\widehat{\Delta}$ -juncture-states are zero.

We are now ready to state our version of construction of the sought-for Gabella quantum holonomy.

Definition 4.23 (enhanced and modified version of Gabella quantum holonomy). Let (Σ, \mathcal{P}) , \mathfrak{S} , Δ and $\widehat{\Delta}$ be as in Def.4.8. Let $\omega \in \mathbb{C}^*$, and let $q = \omega^4$. Let (K, s) be a stated oriented VH-tangle in $\mathfrak{S} \times (-1, 1)$.

Isotope (K, s) into a stated oriented VH-tangle (K', s') in $\mathfrak{S} \times (-1, 1)$ in a good position with respect to $\widehat{\Delta}$ (Lem.3.23) through a VH-isotopy.

For each Gabella lift \widetilde{K}^J in $\widetilde{\Sigma}_\Delta \times (-1, 1)$ of (K', s') associated to a $\widehat{\Delta}$ -juncture-state $J : \{\widehat{\Delta}\text{-junctures of } K'\} \rightarrow \{+, -\}$ of K' that restricts to s' at $\partial K'$, we define a monomial $\widehat{Z}_{\widetilde{K}^J}$ in the (square-root) generators $\widehat{Z}_1, \dots, \widehat{Z}_n$ of the Chekhov-Fock algebra $\mathcal{T}_\Delta^\omega$ (Def.3.9) and a coefficient $\underline{\Omega}(\widetilde{K}^J; \omega) \in \mathbb{Z}[\omega, \omega^{-1}]$ as follows.

(G1) (the monomial part) For each edge e of the original triangulation Δ , denote again by e one of any of the two edges e and e' in $\widehat{\Delta}$ corresponding to e . Let

$$b_e(J) \in \mathbb{Z}$$

be the sum of signs of all $\widehat{\Delta}$ -junctures of K' over e assigned by the $\widehat{\Delta}$ -juncture-state J , where $+$ and $-$ are thought of as 1 and -1 respectively. Now let

$$\widehat{Z}_{\widetilde{K}^J} := \left[\prod_{e \in \Delta} \widehat{Z}_e^{b_e(J)} \right]_{\text{Weyl}} \in \mathcal{T}_\Delta^\omega,$$

where $[\sim]_{\text{Weyl}}$ is the Weyl-ordered monomial defined in Def.2.18.

(G2) (the q -power coefficient part) To each of the crossings of \widetilde{K}^J (as defined at the end of §4.1), associate the sign $+1$ if it is of type $(+)$ in Fig.8, and -1 if it is of type $(-)$ in Fig.8. Let

$$\text{wr}_{\widehat{\Delta}}(\widetilde{K}^J) \in \mathbb{Z}$$

be the net sum of all signs of crossings over all triangles of $\widehat{\Delta}$, i.e. the usual writhe of the tangle \widetilde{K}^J considered only over triangles of $\widehat{\Delta}$.

(G3) (the biangle factor; ‘R-matrix’ and cup/cap) For each $i = 1, \dots, n$, let $L_i := K' \cap ((\Sigma_i \setminus \mathcal{P}_i) \times (-1, 1))$ be the part of K' over the biangle $B_i = (\Sigma_i, \mathcal{P}_i)$ of $\widehat{\Delta}$ corresponding to the edge e_i of Δ . Then $(L_i, J|_{\partial L_i})$ is a stated oriented VH-tangle in $(\Sigma_i \setminus \mathcal{P}_i) \times (-1, 1)$. Consider its value under the biangle quantum holonomy map $\text{TrHol}_{B_i}^\omega([L_i, J|_{\partial L_i}]) \in \mathbb{C}$.

Let

$$(4.7) \quad \underline{\Omega}(\widetilde{K}^J; \omega) := \left(\prod_{i=1}^n \text{TrHol}_{B_i}^\omega([L_i, J|_{\partial L_i}]) \right) \cdot q^{-\text{wr}_{\widehat{\Delta}}(\widetilde{K}^J)} \in \mathbb{Z}[\omega, \omega^{-1}].$$

Define the (enhanced) Gabella quantum holonomy for the stated oriented VH-tangle (K, s) in $\mathfrak{S} \times (-1, 1)$ by the formula

$$(4.8) \quad \text{TrHol}_{\Delta}^\omega(K, s) := \sum_{\widetilde{K}^J} \underline{\Omega}(\widetilde{K}^J; \omega) \widehat{Z}_{\widetilde{K}^J} \in \mathcal{T}_{\Delta}^\omega$$

where the sum is over all isotopy classes of admissible Gabella lifts \widetilde{K}^J of (K', s') (Def.4.8), i.e. over all admissible $\widehat{\Delta}$ -junction-states J of K' that restricts to s' at $\partial K'$.

One might wonder why one can choose any one of e and e' in (G1) above. We claim that for any $\widehat{\Delta}$ -junction-state J of K' for which the coefficient $\underline{\Omega}(\widetilde{K}^J; \omega)$ is nonzero, we have $b_e(J) = b_{e'}(J)$; this is an easy consequence of the charge conservation property (Lem.4.18(2)) of the biangle quantum holonomy. The ideas of (G1) and (G2) already appeared in [GLM15] for a connected non-closed tangle that is given the always-going-up elevations, and as mentioned already, the main contribution of Gabella [G17] is the biangle factor (G3), which enables one to define the quantum holonomy for closed tangles. Meanwhile, as mentioned in §4.2, the biangle quantum holonomy was only partially dealt with in [G17], and the independence of the value of $\text{TrHol}_{\Delta}^\omega(K, s)$ (even in case $K = \emptyset$) on the choice made in the construction, namely the choice of an isotopy transformation into a tangle K' in a good position, was only partially proved in [G17]. This independence was made as a conjecture in the earlier version of the present paper [KSv1], and will be obtained as a consequence of the main result of the present version of the present paper, in a fully general case of a stated oriented VH-tangle.

In the meantime, one actual difference between Gabella’s original construction [G17] and ours is the monomial part (G1); Gabella uses different normalization. Namely, denote by $|b_e|(K)$ be the number of all junctions of K on e . Then, in place of our monomial $\widehat{Z}_{\widetilde{K}^J}$, Gabella uses

$$(4.9) \quad \widehat{X}_{\widetilde{K}^J} := \left[\prod_{e \in T} \widehat{X}_e^{\frac{1}{2}(b_e(J) + |b_e|(K))} \right]_{\text{Weyl}} \in \mathcal{T}_{\Delta}^q.$$

Gabella’s choice also makes the final result to enjoy many properties, and has an advantage of avoiding the square-root variables ($\cdot: \frac{1}{2}(b_e(J) + |b_e|(K)) \in \mathbb{Z}$) and making the lowest term to be 1. Later, we will discuss why we had to modify as we did.

As already mentioned, the Gabella’s original construction is for a simple loop in the surface \mathfrak{S} .

Definition 4.24. Let Δ be a triangulation of a triangulable generalized marked surface (Σ, \mathcal{P}) , and let γ be an oriented simple loop in $\mathfrak{S} = \Sigma \setminus \mathcal{P}$.

Let K_γ be the oriented tangle in $\mathfrak{S} \times (-1, 1)$ obtained as a constant-elevation lift of γ , as in Def.3.17. Define the Gabella quantum holonomy of the oriented simple loop γ in the surface \mathfrak{S} as

$$(4.10) \quad \text{TrHol}_{\Delta}^\omega(\gamma) := \text{TrHol}_{\Delta}^\omega(K_\gamma, \emptyset) \in \mathcal{T}_{\Delta}^\omega.$$

Some of the important properties of $\text{TrHol}_{\Delta}^\omega(\gamma)$ proved by Gabella in [G17] read as follows, when translated to our normalization:

Proposition 4.25 (properties of Gabella quantum holonomy; [G17]). Let (Σ, \mathcal{P}) , \mathfrak{S} , Δ , γ and $K = K_\gamma$ be as in Def.4.24. Suppose further that γ is not a contractible loop in \mathfrak{S} .

- (1) When $\omega = 1$, $\text{TrHol}_{\Delta}^1(\gamma)$ recovers the (absolute value of the) usual trace of holonomy function $\mathbb{I}(\gamma)$ on $\mathcal{X}^+(\Sigma, \mathcal{P})$ along γ ;

- (2) In terms of the partial ordering on the monomials $\widehat{Z}_{\widetilde{K}^J} \in \mathcal{T}_\Delta^\omega$ appearing in the right hand side of the sum formula eq.(4.8) for $\text{TrHol}_\Delta^\omega(\gamma) = \text{TrHol}_\Delta^\omega(K, \emptyset)$ induced by the powers of the generators $\widehat{Z}_1, \dots, \widehat{Z}_n$, there is a unique Gabella lift \widetilde{K}^J giving the highest term, for which the power a_e of each generator \widehat{Z}_e in the monomial $\widehat{Z}_{\widetilde{K}^J}$, associated to the edge e of Δ , equals the intersection number $a_e(\gamma)$ defined in Lem.2.8 (i.e. 2 times the Fock-Goncharov tropical \mathcal{A} -coordinate for the lamination γ), and the coefficient $\underline{\Omega}(\widetilde{K}^J; \omega)$ is 1;
- (3) For each \widetilde{K}^J appearing in eq.(4.8), the coefficient $\underline{\Omega}(\widetilde{K}^J; \omega)$ belongs to $\mathbb{Z}_{\geq 0}[q, q^{-1}]$, i.e. a positive integral Laurent polynomial in $q = \omega^4$;
- (4) For each \widetilde{K}^J appearing in eq.(4.8), the coefficient $\underline{\Omega}(\widetilde{K}^J; \omega)$ is $*$ -invariant, in the sense that $\underline{\Omega}(\widetilde{K}^J; \omega) = \underline{\Omega}(\widetilde{K}^J; \omega^{-1})$, or that it is invariant under the exchange $q \leftrightarrow q^{-1}$.

One remark is that the proof given in [G17, §6.4] of the positivity property of part (3) is not quite sufficient; we note that this positivity will follow from the main result of our paper along with [CKKO20].

5. EQUALITY OF THE TWO CONSTRUCTIONS

5.1. Statement of the main theorem. For a stated oriented VH-tangle (K, s) in $\mathfrak{S} \times (-1, 1)$, we investigated the Bonahon-Wong quantum trace $\text{Tr}_\Delta^\omega([K, s]) \in \mathcal{T}_\Delta^\omega$ (by viewing (K, s) as a stated oriented V-tangle) and Gabella's quantum holonomy $\text{TrHol}_\Delta^\omega([K, s]) \in \mathcal{T}_\Delta^\omega$. They have several properties in common, which naturally leads to the question of whether they are equal, as mentioned in [G17] for the case when K is closed and connected. This equality is the main theorem of the present paper. In order to formulate the result in full generality incorporating possibly non-closed tangles, we first need to introduce the following.

Definition 5.1 (signed order correction amount). *Let (Σ, \mathcal{P}) be a generalized marked surface, not necessarily triangulable. Let $\mathfrak{S} = \Sigma \setminus \mathcal{P}$. Let e be a \mathcal{P} -arc (Def.2.3) of (Σ, \mathcal{P}) , with an orientation chosen on e .*

• *Let Z be an ordered finite subset of e , and $s_0 : Z \rightarrow \{+, -\}$ be a state of Z . Define the signed order correction amount of the stated ordered subset (Z, s_0) on the oriented arc e as the integer*

$$\mathcal{C}(e; Z, s_0) := \sum_{x, y \in Z \mid x < y} \text{sgn}(e; \overrightarrow{xy}) s_0(x) s_0(y) \in \mathbb{Z},$$

where each sign \pm is regarded as the number ± 1 , and

$$\text{sgn}(e; \overrightarrow{xy}) := \begin{cases} 1, & \text{if the direction from } x \text{ to } y \text{ matches the given orientation on } e, \\ -1, & \text{otherwise.} \end{cases}$$

When $Z = \emptyset$, we set $\mathcal{C}(e; \emptyset, \emptyset) = 0$.

• *When Z is a finite subset of the thickening $e \times (-1, 1)$ of e equipped with a state $s_0 : Z \rightarrow \{+, -\}$, such that the restrictions to Z of the projection maps $e \times (-1, 1) \rightarrow e$ and $e \times (-1, 1) \rightarrow (-1, 1)$ are injective, we define $\mathcal{C}(e; Z, s_0)$ using the projection of (Z, s_0) onto e , with the ordering on the projection of Z coming from the elevations of the elements of Z , i.e. the vertical ordering.*

• *When e is a boundary arc and no orientation is specified, we use the boundary-orientation (Def.2.1) coming from the surface orientation on Σ .*

• *If (K, s) is a stated VH-tangle in $\mathfrak{S} \times (-1, 1)$, the boundary signed order correction amount of (K, s) is defined as*

$$\partial\mathcal{C}_{(\Sigma, \mathcal{P})}(K, s) := \sum_{\text{boundary arcs } b \text{ of } (\Sigma, \mathcal{P})} \mathcal{C}(b; \partial K \cap (b \times (-1, 1)), s_0|_{\partial K \cap (b \times (-1, 1))})$$

where we always use the boundary-orientation on b 's even if (Σ, \mathcal{P}) is a directed biangle.

We now state the main theorem.

Theorem 5.2 (main theorem). *Let (Σ, \mathcal{P}) be a triangulable generalized marked surface, and Δ a triangulation of (Σ, \mathcal{P}) ; let $\mathfrak{S} = \Sigma \setminus \mathcal{P}$. Let $\omega \in \mathbb{C}^*$, $q = \omega^4$ and $A = \omega^{-2}$. Let (K, s) be a stated oriented VH-tangle in $\mathfrak{S} \times (-1, 1)$. Then the Bonahon-Wong quantum trace $\text{Tr}_\Delta^\omega([K, s]) \in \mathcal{Z}_\Delta^\omega \subset \mathcal{T}_\Delta^\omega$,*

constructed in §3 for the stated skein $[K, s] \in \mathcal{S}_s^A(\Sigma, \mathcal{P})$ is related to the enhanced Gabella quantum holonomy $\text{TrHol}_\Delta^\omega(K, s) \in \mathcal{T}_\Delta^\omega$ constructed in §4 as

$$(5.1) \quad \text{Tr}_\Delta^\omega([K, s]) = \omega^{2\text{wr}(K)} \omega^{\partial\mathcal{C}_{(\Sigma, \mathcal{P})}(K, s)} \text{TrHol}_\Delta^\omega(K, s),$$

where the writhe $\text{wr}(K) \in \mathbb{Z}$ of the VH-tangle K is defined as

$$\text{wr}(K) := \#(\text{crossings of type } (+) \text{ in Fig.8}) - \#(\text{crossings of type } (-) \text{ in Fig.8}),$$

counted over Σ (not over the branched double cover $\tilde{\Sigma}_\Delta$).

In particular, this equality holds for a constant-elevation lift $K = K_\gamma$ of an oriented simple loop γ in \mathfrak{S} (Def.3.17), where $\text{wr}(K) = 0$, $\partial K = \emptyset$, $s = \emptyset$ and $\partial\mathcal{C}_{(\Sigma, \mathcal{P})}(K, \emptyset) = 0$. Hence, Allegretti-Kim's solution $\widehat{\mathbb{I}}_\Delta^\omega(\gamma)$ in eq.(3.6) to the quantum ordering problem for the classical function $\mathbb{I}(\gamma)$ addressed in §2.4 coincides with Gabella's solution $\text{TrHol}_\Delta^\omega(\gamma)$ in eq.(4.10).

This theorem has some immediate consequences on the Gabella quantum holonomy $\text{TrHol}_\Delta^\omega(K, s)$, following from the corresponding properties of the Bonahon-Wong quantum trace $\text{Tr}_\Delta^\omega([K, s])$ and some obvious observations about $\text{wr}(K)$ and $\partial\mathcal{C}_{(\Sigma, \mathcal{P})}(K, s)$: namely, we obtain a proof of Prop.4.25, especially the positivity in part (3), and the following corollaries.

Corollary 5.3 (isotopy invariance). *The enhanced Gabella quantum holonomy $\text{TrHol}_\Delta^\omega(K, s)$ defined in Def.4.23 depends only on the VH-isotopy class of a stated oriented VH-tangle (K, s) .*

Corollary 5.4 (mutation compatibility). *Let (Σ, \mathcal{P}) , Δ , $\widehat{\Delta}$ be as in Def.3.22; let $\mathfrak{S} = \Sigma \setminus \mathcal{P}$. Let $\omega \in \mathbb{C}^*$, $q = \omega^4$. Let Δ' be another triangulation of (Σ, \mathcal{P}) , and $\widehat{\Delta}'$ be a corresponding split ideal triangulation of Δ' .*

Let (K, s) be an oriented stated VH-tangle in $\mathfrak{S} \times (-1, 1)$. Then we have

$$\text{TrHol}_\Delta^\omega(K, s) = \Theta_{\Delta\Delta'}^\omega(\text{TrHol}_{\Delta'}^\omega(K, s)),$$

where $\Theta_{\Delta\Delta'}^\omega$ is the square-root quantum coordinate change map as in Prop.3.10. In particular, $\text{TrHol}_\Delta^\omega(K, s) \in \mathcal{Z}_\Delta^\omega$ and $\text{TrHol}_{\Delta'}^\omega(K, s) \in \mathcal{Z}_{\Delta'}^\omega$ (Def.3.9).

As a consequence, we also obtain $\text{TrHol}_\Delta^\omega(K, s) \in \mathbf{L}_\Delta^\omega$, where \mathbf{L}_Δ^ω is defined as in eq.(3.2).

The main theorem and these corollaries hold for the Gabella quantum holonomy $\text{TrHol}_\Delta^\omega(K, s)$ defined using the normalization Def.4.23.(G1) as in our present paper, but not for Gabella's original construction [G17] which uses the normalization in eq.(4.9).

We note that, in the earlier version [KSv1] of the present paper, the main theorem was proved only for the case when the tangle K is closed, connected, and has no crossing, because the biangle quantum holonomy was only partially dealt with. In the present version, through the contribution of the newly participating second author, we now have a complete treatment of the biangle quantum holonomy, hence the main theorem in full generality. The new idea added in this version also made the proofs shorter and more direct.

Let's now start proving the main theorem. Let (K, s) be any stated oriented VH-tangle in $\mathfrak{S} \times (-1, 1)$. Isotope it to a stated oriented VH-tangle (K', s') in a good position with respect to $\widehat{\Delta}$, through a VH-isotopy. Then both sides of sought-for eq.(5.1) are expressed as sums over admissible $\widehat{\Delta}$ -juncture-states J of K' restricting to s' at ∂K ; see Prop.3.25 ad Def.4.23. It is enough to show that the summands of the two sides for each J coincide, i.e. to show the term-by-term equality

$$(5.2) \quad \text{BW}_\Delta^\omega(K'; J) = \omega^{2\text{wr}(K)} \omega^{\partial\mathcal{C}_{(\Sigma, \mathcal{P})}(K, s)} \overline{\Omega}(\widetilde{K}^J; \omega) \widehat{Z}_{\widetilde{K}^J}$$

for each J . The rest of the present section is devoted to a proof of the term-by-term equality, eq.(5.2); for convenience, from now on we may identify K with K' :

$$K = K'$$

Throughout the entire section, we will reserve the symbols K , K' and s as such.




FIGURE 9. Pair of tangle segments over triangle (arrows in boundary are clockwise)

5.2. Bonahon-Wong triangle factors. We further break down the Bonahon-Wong summand $\text{BW}_{\widehat{\Delta}}^{\omega}(K'; J) = \text{BW}_{\widehat{\Delta}}^{\omega}(K; J)$ in the left hand side of the sought-for eq.(5.2) into the product of the biangle factors $\text{Tr}_{B_i}^{\omega}([L_i, J|_{\partial L_i}])$ and the triangle factors $\text{Tr}_{\widehat{t}_j}^{\omega}([K_j, J|_{\partial K_j}])$, as in its very definition in eq.(3.11). In the present subsection we focus on the triangle factors.

Let (Σ, \mathcal{P}) , \mathfrak{S} , Δ , $\widehat{\Delta}$, e_i , e'_i , B_i , t_j and \widehat{t}_j be as in Def.3.22. Let $\omega \in \mathbb{C}^*$ and $q = \omega^4$. As done in Prop.3.25, we let $K_j := K \cap (\widehat{t}_j \times (-1, 1))$ be the part of K over the triangle \widehat{t}_j of $\widehat{\Delta}$, and let $k_{j,1}, \dots, k_{j,l_j}$ the components of K_j , i.e. the triangle segments of K over \widehat{t}_j , in order of increasing elevation. The triangle factor for this triangle \widehat{t}_j is as written in eq.(3.10):

$$\text{Tr}_{\widehat{t}_j}^{\omega}([K_j, J|_{\partial K_j}]) = \text{Tr}_{\widehat{t}_j}^{\omega}([k_{j,1}, J|_{\partial k_{j,1}}]) \text{Tr}_{\widehat{t}_j}^{\omega}([k_{j,2}, J|_{\partial k_{j,2}}]) \cdots \text{Tr}_{\widehat{t}_j}^{\omega}([k_{j,l_j}, J|_{\partial k_{j,l_j}}]) \in \mathcal{T}_{\widehat{t}_j}^{\omega}.$$

We will now rewrite the right hand side as the Weyl-ordered monomial times some integer power of ω , which we refer to as the *deviation*.

Definition 5.5. *The deviation of the Bonahon-Wong triangle factor for triangle \widehat{t}_j from the Weyl-ordering, which is associated to the stated V -tangle $(K_j, J|_{\partial K_j})$ in the thickening of \widehat{t}_j , is the unique integer $\text{dev}_{\widehat{t}_j}(K_j, J|_{\partial K_j})$ such that*

$$\text{Tr}_{\widehat{t}_j}^{\omega}([K_j, J|_{\partial K_j}]) = \omega^{\text{dev}_{\widehat{t}_j}(K_j, J|_{\partial K_j})} \left[\text{Tr}_{\widehat{t}_j}^{\omega}([k_{j,1}, J|_{\partial k_{j,1}}]) \cdots \text{Tr}_{\widehat{t}_j}^{\omega}([k_{j,l_j}, J|_{\partial k_{j,l_j}}]) \right]_{\text{Weyl}}.$$

If J is not admissible, both sides of the above equation are zero, so the deviation would not be uniquely determined; but we only deal with admissible J 's. We could apply the definition of deviation to more general stated V -tangles satisfying certain conditions, e.g. (GP2)–(GP3) of Def.3.22. For example, for the stated V -tangle consisting of two of the components of $(K_j, J|_{\partial K_j})$, whenever $1 \leq r < u \leq l_j$ we have

$$\text{Tr}_{\widehat{t}_j}^{\omega}([k_{j,r}, J|_{\partial k_{j,r}}]) \text{Tr}_{\widehat{t}_j}^{\omega}([k_{j,u}, J|_{\partial k_{j,u}}]) = \omega^{\text{dev}_{\widehat{t}_j}(k_{j,r} \cup k_{j,u}, J|_{\partial k_{j,r} \cup \partial k_{j,u}})} \left[\text{Tr}_{\widehat{t}_j}^{\omega}([k_{j,r}, J|_{\partial k_{j,r}}]) \text{Tr}_{\widehat{t}_j}^{\omega}([k_{j,u}, J|_{\partial k_{j,u}}]) \right].$$

It is straightforward to prove the following well-known observation, which enables us to express the deviation for a triangle as a sum of deviations over pairs of tangle segments in the triangle:

$$(5.3) \quad \text{dev}_{\widehat{t}_j}(K_j, J|_{\partial K_j}) = \sum_{1 \leq r < u \leq l_j} \text{dev}_{\widehat{t}_j}(k_{j,r} \cup k_{j,u}, J|_{\partial k_{j,r} \cup \partial k_{j,u}}).$$

We establish a computational lemma for each summand of the right-hand-side of eq.(5.3).

Lemma 5.6 (deviation for a pair of stated tangle segments in a triangle). *If $(k_1 \cup k_2, s_0)$ is a stated V -tangle in the thickening of a non-self-folded triangle \widehat{t} consisting of two tangle segments k_1 and k_2 as in the cases (1) or (2) of Fig.9, and if we denote $\varepsilon_1 = s_0(x_1)$, $\varepsilon_2 = s_0(x_2)$, $\varepsilon_3 = s_0(x_3)$ and $\varepsilon_4 = s_0(x_4)$, then we have*

$$\text{dev}_{\widehat{t}}(k_1 \cup k_2, s_0) = \begin{cases} \varepsilon_1 \varepsilon_4 - \varepsilon_2 \varepsilon_3, & \text{in case (1),} \\ \varepsilon_1 \varepsilon_3 - \varepsilon_1 \varepsilon_4 + \varepsilon_2 \varepsilon_4 & \text{in case (2).} \end{cases}$$

Proof. Follows from straightforward computation. \blacksquare

5.3. Gabella triangle factors. Now we turn to the triangle factors of the term for Gabella quantum holonomy, i.e. the right hand side of eq.(5.2). Recall from (4.7) that $\underline{\Omega}(\widetilde{K}^J; \omega)$ is the product of the biangle quantum holonomy factors for biangles and a power of q to (minus of) the writhe of the Gabella lift \widetilde{K}^J counted over triangles of $\widehat{\Delta}$. One must be careful that this writhe is considered over the branched double cover surface $\widehat{\Sigma}_{\Delta}$, and not over Σ . Writhe is a net sum of signs of all crossings, where each crossing

is formed by two segments living over a same triangle of $\widehat{\Delta}$. Thus we can write this writhe $\text{wr}_{\widehat{\Delta}}(\widetilde{K}^J)$ as a double sum, where the outer sum is over triangles of $\widehat{\Delta}$ and the inner sum is over pairs of segments living over each triangle. For triangle \widehat{t}_j of $\widehat{\Delta}$, denote by \widetilde{K}_j^J the part of \widetilde{K}^J over \widehat{t}_j ; more precisely, the subset of \widetilde{K}^J of points whose images under the composition $\widetilde{\Sigma}_{\Delta} \times (-1, 1) \rightarrow \Sigma \times (-1, 1) \rightarrow \Sigma$ lie in \widehat{t}_j . Define $\text{wr}_{\widehat{t}_j}(\widetilde{K}_j^J)$ be the writhe of \widetilde{K}_j^J , i.e. the net sum of signs of the crossings (see Def.4.23(G2)) of \widetilde{K}_j^J . Then

$$\text{wr}_{\widehat{\Delta}}(\widetilde{K}^J) = \sum_{j=1}^m \text{wr}_{\widehat{t}_j}(\widetilde{K}_j^J)$$

Note that \widetilde{K}_j^J is a Gabella lift of the stated oriented VH-tangle K_j living in the thickening of the triangle \widehat{t}_j . Denote the components of \widetilde{K}_j^J as $\widetilde{k}_{j,1}, \dots, \widetilde{k}_{j,l_j}$. Then one can observe that each of these is a Gabella lift of the corresponding one among the components of K_j which are stated oriented tangle segments $(k_{j,1}, J|_{\partial k_{j,1}}), \dots, (k_{j,l_j}, J|_{\partial k_{j,l_j}})$. That is, $\widetilde{k}_{j,r}$ is a Gabella lift of the horizontal triangle segment $k_{j,r}$ with the state coinciding with $J|_{\partial k_{j,r}}$ (Def.4.7). One observes

$$(5.4) \quad \text{wr}_{\widehat{t}_j}(\widetilde{K}_j^J) = \sum_{1 \leq r < u \leq l_j} \text{wr}_{\widehat{t}_j}(\widetilde{k}_{j,r} \cup \widetilde{k}_{j,u}),$$

thus indeed we expressed $\text{wr}_{\widehat{\Delta}}(\widetilde{K}^J)$ as a double sum. Now we establish a computational lemma for the innermost summand.

Lemma 5.7 (writhe for a pair of stated oriented tangle segments in a triangle). *Let \widehat{t} be a non-self-folded triangle, viewed as a generalized marked surface (Σ', \mathcal{P}') , with the unique ideal triangulation Δ' . Let $\mathfrak{S}' = \Sigma' \setminus \mathcal{P}'$. Let $(k_1 \cup k_2, s_0)$ be a stated oriented VH-tangle in $\mathfrak{S}' \times (-1, 1)$ as in the case (1) or (2) of Fig.9, with arbitrary orientations on k_1 and k_2 . Let $\widetilde{k}_1 \cup \widetilde{k}_2$ be a Gabella lift in $\widetilde{\Sigma}'_{\Delta'} \times (-1, 1)$ of $(k_1 \cup k_2, s_0)$ (in the sense of Def.4.8), where each of \widetilde{k}_1 and \widetilde{k}_2 is a Gabella lift of $(k_1, s_0|_{\partial k_1})$ and $(k_2, s_0|_{\partial k_2})$ respectively (in the sense of Def.4.7 with the prescribed states).*

If we denote $\varepsilon_1 = s_0(x_1)$, $\varepsilon_2 = s_0(x_2)$, $\varepsilon_3 = s_0(x_3)$ and $\varepsilon_4 = s_0(x_4)$, then the writhe of the Gabella lift $\widetilde{k}_1 \cup \widetilde{k}_2$ is given by:

$$\text{wr}_{\widehat{t}}(\widetilde{k}_1 \cup \widetilde{k}_2) = \begin{cases} -\frac{1}{4}(\varepsilon_1 - \varepsilon_2)(\varepsilon_3 + \varepsilon_4), & \text{in case (1),} \\ -\frac{1}{4}(\varepsilon_1 - \varepsilon_2)(\varepsilon_3 - \varepsilon_4), & \text{in case (2),} \end{cases}$$

where each sign \pm is understood as the number ± 1 .

Remark 5.8. *This is another occurrence of the phenomenon of orientation insensitivity.*

Proof. Note that the Gabella lift $\widetilde{k}_1 \cup \widetilde{k}_2$ is uniquely determined up to equivalence, in this situation; any Gabella lift gives the same answer for the writhe, because writhe is well-defined up to (VH-)isotopy. Consider the case (1) of Fig.9, where both segments k_1 and k_2 live in the same corner; with respect to the vertex of this corner, k_1 is located ‘inner’ and k_2 ‘outer’. If $\varepsilon_1 = \varepsilon_2$ then the Gabella lift \widetilde{k}_1 of the inner segment bounds a corner region of the triangle not containing the branch point (see Fig.5), hence can be isotoped so that it doesn’t meet \widetilde{k}_2 . If $\varepsilon_3 \neq \varepsilon_4$ the the Gabella lift \widetilde{k}_2 of the outer segment bounds a corner region of the triangle containing the branch point (see Fig.5), hence can be isotoped so that it doesn’t meet \widetilde{k}_1 . So, in these cases, we have $\text{wr}_{\widehat{t}}(\widetilde{k}_1 \cup \widetilde{k}_2) = 0$; meanwhile, note that $-\frac{1}{4}(\varepsilon_1 - \varepsilon_2)(\varepsilon_3 + \varepsilon_4)$ equals zero in these cases, so we get the sought-for equality. Now assume that both $\varepsilon_1 \neq \varepsilon_2$ and $\varepsilon_3 = \varepsilon_4$ hold; there are four such possibilities for signs. The upper half of Fig.10 presents the diagram of $\widetilde{k}_1 \cup \widetilde{k}_2$ for each of these four possibilities, under particular choice of orientations of k_1 and k_2 ; one can easily verify the equality in these cases. For each case in the upper half of Fig.10, note that the projections of \widetilde{k}_1 and \widetilde{k}_2 in \widehat{t} meet at two points in Σ , but only one of which is a crossing point over the branched double cover $\widetilde{\Sigma}_{\Delta}$ of $\widetilde{k}_1 \cup \widetilde{k}_2$. If one changes the orientation of either one of k_1 and k_2 , say k_i , then the projections of \widetilde{k}_1 and \widetilde{k}_2 in \widehat{t} are same as before, with the orientation of k'_i reversed from before. Then the diagram of each of \widetilde{k}_1 and \widetilde{k}_2 stay the same (though the diagram of $\widetilde{k}_1 \cup \widetilde{k}_2$ is changed), with the orientation of \widetilde{k}_i reversed, and the sheet numbers of points of \widetilde{k}_i changed from before. So, out of the two intersections of the projections of \widetilde{k}_1 and \widetilde{k}_2 in \widehat{t} , the one that used to be the crossing of $\widetilde{k}_1 \cup \widetilde{k}_2$ is not a crossing anymore, and the remaining one now becomes the crossing point in $\widetilde{\Sigma}_{\Delta}$. However,




FIGURE 10. Diagrams for a Gabella lift $\tilde{k}_1 \cup \tilde{k}_2$ in $\tilde{\Sigma}_{\Delta}$ of union of two stated oriented VH-tangles in a triangle \tilde{t} , for computation of writhe

one easily verifies that the sign of the crossing is same as before. Similar argument holds whenever one changes the orientation of both k_1 and k_2 .

Now consider the case (2) of Fig.9. If $\varepsilon_1 = \varepsilon_2$ or $\varepsilon_3 = \varepsilon_4$ holds, it is easy to see that both sides of the equation in the statement are zero. When both $\varepsilon_1 \neq \varepsilon_2$ and $\varepsilon_3 \neq \varepsilon_4$ hold, one can check the equality case by case, for each of the four possibilities for the signs; see the lower half of Fig.10, drawn for particular choices of orientations. Similarly as in the case (1), one easily observes that the equality holds also for the other choices of orientations. ■

Using the computational lemmas developed so far, we now arrive at the relationship between the triangle factors of the Bonahon-Wong term and the Gabella term. In fact, they are not exactly equal, but equal only up to powers of ω to the signed order correction amounts defined in Def.5.1, which is due to the fact that Bonahon-Wong quantum trace is defined for V-tangles up to V-isotopy, but writhe (and hence Gabella quantum holonomy) is defined for VH-tangles up to VH-isotopy. The precise statement is as follows, and is the main technical lemma of the proof of our main theorem.

Lemma 5.9 (equality of the triangle factors). *One has*

$$\text{dev}_{\tilde{t}_j}(K_j, J|_{\partial K_j}) = -4wr_{\tilde{t}_j}(\tilde{K}_j^J) + 2wr(K_j) + \partial\mathcal{C}_{\tilde{t}_j}(K_j, J|_{\partial K_j})$$

Proof. We first re-write the boundary signed order correction amount as the sum over pairs of tangle segments $k_{j,1}, \dots, k_{j,l_j}$ of K_j :

$$(5.5) \quad \partial\mathcal{C}_{\tilde{t}_j}(K_j, J|_{\partial K_j}) = \sum_{1 \leq r < u \leq l_j} \partial\mathcal{C}_{\tilde{t}_j}(k_{j,r} \cup k_{j,u}, J|_{\partial k_{j,r} \cup \partial k_{j,u}});$$

this is easily proved by looking at the contributions at each boundary arc of \tilde{t}_j .

Let $1 \leq r < u \leq l_j$. In view of the sum expression for the deviation obtained in eq.(5.3), and of

$$wr(K_j) = \sum_{1 \leq r < u \leq l_j} wr(k_{j,r} \cup k_{j,u}),$$

we now investigate the number

$$(*)_{j;r,u} := \text{dev}_{\tilde{t}_j}(k_{j,r} \cup k_{j,u}, J|_{\partial k_{j,r} \cup \partial k_{j,u}}) + 4wr_{\tilde{t}_j}(\tilde{k}_{j,r} \cup \tilde{k}_{j,u}) - 2wr(k_{j,r} \cup k_{j,u}),$$

and will show that it equals the (r, u) -th summand in the right hand side of eq.(5.5). We apply Lem.5.6 to $(k_{j,r} \cup k_{j,u}, J|_{\partial k_{j,r} \cup \partial k_{j,u}})$, for which we also apply Lem.5.7 with the Gabella lift $\tilde{k}_{j,r} \cup \tilde{k}_{j,u}$.

If $k_{j,r}$ and $k_{j,u}$ are like k_1 and k_2 of the case (1) of Fig.9, we have

$$\begin{aligned}
(*)_{j;r,u} &= \underbrace{(J(x_1)J(x_4) - J(x_2)J(x_3))}_{\text{Lem.5.6}} - \underbrace{(J(x_1) - J(x_2))(J(x_3) + J(x_4))}_{\text{Lem.5.7}} - \underbrace{2\text{wr}(k_{j,r} \cup k_{j,u})}_{=0} \\
&= (-J(x_1)J(x_3) + J(x_2)J(x_4)) \\
&= \text{sgn}(e; \overrightarrow{x_1x_3})J(x_1)J(x_3) + \text{sgn}(e'; \overrightarrow{x_2x_4})J(x_2)J(x_4),
\end{aligned}$$

where e is the side of \widehat{t}_j containing x_1 and x_3 , while e' is the side containing x_2 and x_4 ; note $x_1 < x_3$ and $x_2 < x_4$ in the ordering coming from elevations, because $k_{j,r}$ has lower elevation than $k_{j,u}$. Therefore

$$(5.6) \quad (*)_{j;r,u} = \partial\mathcal{C}_{\widehat{t}_j}(k_{j,r} \cup k_{j,u}, J|_{\partial k_{j,r} \cup \partial k_{j,u}})$$

holds in this case, as desired.

Suppose now that $k_{j,r}$ and $k_{j,u}$ are like k_1 and k_2 of the case (3) of Fig.9. As tangles, this case can be obtained from case (1) by a V-isotopy that slides the endpoints x_1 of k_1 and x_3 of k_2 . After this sliding (or boundary exchange move (M2) in §3.1), note that $\text{dev}_{\widehat{t}_j}(k_{j,r} \cup k_{j,u}, J|_{\partial k_{j,r} \cup \partial k_{j,u}})$ does not change, and the boundary signed order correction amount $\partial\mathcal{C}_{\widehat{t}_j}(k_{j,r} \cup k_{j,u}, J|_{\partial k_{j,r} \cup \partial k_{j,u}})$ increase by 2 if $J(x_1) = J(x_3)$ and decrease by 2 if $J(x_1) \neq J(x_3)$. Suppose that the orientations on k_1 and k_2 are parallel, e.g. going from x_1 to x_2 and x_3 to x_4 . After sliding, $-2\text{wr}(k_{j,r} \cup k_{j,u})$ decreases by 2, while $4\text{wr}_{\widehat{t}_j}(\widetilde{k}_{j,r} \cup \widetilde{k}_{j,u})$ increase by 4 if $J(x_1) = J(x_3)$ (because the parts of the two segments near the boundary edge are living in the same sheet) and stays the same if $J(x_1) \neq J(x_3)$ (because the parts of the two segments near the boundary edge are living in different sheets). Thus eq.(5.6) holds. Now suppose that the orientations of k_1 and k_2 are not parallel; e.g. suppose now k_2 is going from x_4 to x_3 . After the sliding move from case (1), $-2\text{wr}(k_{j,r} \cup k_{j,u})$ increases by 2, while $4\text{wr}_{\widehat{t}_j}(\widetilde{k}_{j,r} \cup \widetilde{k}_{j,u})$ stays the same if $J(x_1) = J(x_3)$ and decrease by 4 if $J(x_1) \neq J(x_3)$. Thus eq.(5.6) still holds.

If $k_{j,r}$ and $k_{j,u}$ are like k_1 and k_2 of the case (2) of Fig.9, we have

$$\begin{aligned}
(*)_{j;r,u} &= \underbrace{(J(x_1)J(x_3) - J(x_1)J(x_4) + J(x_2)J(x_4))}_{\text{Lem.5.6}} - \underbrace{(J(x_1) - J(x_2))(J(x_3) - J(x_4))}_{\text{Lem.5.7}} - \underbrace{2\text{wr}(k_{j,r} \cup k_{j,u})}_{=0} \\
&= J(x_2)J(x_3) = \text{sgn}(e; \overrightarrow{x_2x_3})J(x_2)J(x_3),
\end{aligned}$$

where e is the side of \widehat{t}_j containing x_2 and x_3 ; note $x_2 < x_3$ in the ordering coming from elevations, because $k_{j,r}$ has lower elevation than $k_{j,u}$. Thus eq.(5.6) holds also in this case.

Suppose now that $k_{j,r}$ and $k_{j,u}$ are like k_1 and k_2 of the case (4) of Fig.9. As tangles, this case can be obtained from case (2) by a V-isotopy that slides the endpoints x_2 of k_1 and x_3 of k_2 . After this sliding (or boundary exchange move), note that $\text{dev}_{\widehat{t}_j}(k_{j,r} \cup k_{j,u}, J|_{\partial k_{j,r} \cup \partial k_{j,u}})$ does not change, and the boundary signed order correction amount $\partial\mathcal{C}_{\widehat{t}_j}(k_{j,r} \cup k_{j,u}, J|_{\partial k_{j,r} \cup \partial k_{j,u}})$ decrease by 2 if $J(x_2) = J(x_3)$ and increase by 2 if $J(x_2) \neq J(x_3)$. Suppose that the orientations on k_1 and k_2 are parallel, e.g. going from x_2 to x_1 and x_3 to x_4 . After sliding, $-2\text{wr}(k_{j,r} \cup k_{j,u})$ increases by 2, while $4\text{wr}_{\widehat{t}_j}(\widetilde{k}_{j,r} \cup \widetilde{k}_{j,u})$ decrease by 4 if $J(x_2) = J(x_3)$ and stays the same if $J(x_2) \neq J(x_3)$. Thus eq.(5.6) holds. Now suppose that the orientations of k_1 and k_2 are not parallel; e.g. suppose now k_1 is going from x_1 to x_2 . After the sliding move from case (2), $-2\text{wr}(k_{j,r} \cup k_{j,u})$ decreases by 2, while $4\text{wr}_{\widehat{t}_j}(\widetilde{k}_{j,r} \cup \widetilde{k}_{j,u})$ stays the same if $J(x_2) = J(x_3)$ and increase by 4 if $J(x_2) \neq J(x_3)$. Thus eq.(5.6) still holds.

When $k_{j,r}$ and $k_{j,u}$ play the roles of k_2 and k_1 of Fig.9 in each of the cases we dealt with so far, eq.(5.6) still holds because of the skew-symmetry of each of the three terms of $(*)_{j;r,u}$, as well as that of the boundary signed order correction amount $\partial\mathcal{C}_{\widehat{t}_j}(k_{j,r} \cup k_{j,u}, J|_{\partial k_{j,r} \cup \partial k_{j,u}})$.

Now, taking the sum of the equality eq.(5.6) over all pairs (r, u) with $1 \leq r < u \leq l_j$, we get the desired result, in view of eq.(5.3), eq.(5.4), and eq.(5.5). \blacksquare

The boundary signed order correction amount will also appear in the relationship between the biangle factors of the Bonahon-Wong term and the Gabella term, cancelling the ones appearing for the triangles.

5.4. Bonahon-Wong biangle factors via $\mathcal{U}_q(\mathfrak{sl}_2)$ Reshetikhin-Turaev operator invariant. Now we move on to the biangle factors. Our approach is via the operator invariants of tangles in a biangle, as we did in §4.2 to investigate Gabella's biangle quantum holonomy. We follow notations of §4.2, some of which we recall now. Let $\vec{B} = (B, \text{dir})$ be a directed biangle (Def.4.9) whose boundary arcs

are denoted by b_{in} and b_{out} , and let $\mathcal{D}(\vec{B})$ be the set of all equivalence classes $[D]$ of boundary-ordered oriented tangle diagrams (Def.4.10). Let the vector space V be still be given as in eq.(4.3), with the ordered basis $\{\xi_+, \xi_-\}$. We claimed in Prop.4.15 that there exists a unique assignment G that assigns to each $[D] \in \mathcal{D}(\vec{B})$ a linear operator $G([D]) : V^{\otimes |\partial_{\text{in}} D|} \rightarrow V^{\otimes |\partial_{\text{out}} D|}$ satisfying certain properties, where $\partial_{\text{in}} D = \partial_{b_{\text{in}}} D$ and $\partial_{\text{out}} D = \partial_{b_{\text{out}}} D$. Then the Gabella biangle quantum holonomy $\text{TrHol}_{\vec{B}}^{\omega}([k, s_0])$ for a stated oriented VH-tangle (k, s_0) in the thickening of \vec{B} was defined as a matrix element of the operator $G([D])$ (Def.4.17), where D is the boundary-ordered tangle diagram of k . We still have to prove Prop.4.15, and also to find a relationship between the Gabella biangle quantum holonomy and the Bonahon-Wong biangle quantum trace dealt with in §3.4. We accomplish these two tasks simultaneously.

The main ingredient of the present section is the Reshetikhin-Turaev operator invariant [RT90] [T16] for oriented tangles in a biangle, associated to the irreducible 2-dimensional representation of the quantum group $\mathcal{U}_q(\mathfrak{sl}_2)$. While suggesting the readers to consult the excellent monograph by Ohtsuki [O02] for details on the theory of operator invariants, we just state and use the known result here.

Proposition 5.10 (Reshetikhin-Turaev operator invariant for 2d irreducible representation of $\mathcal{U}_2(\mathfrak{sl}_2)$; [RT90] [T16] [O02]). *There exists a unique map F assigning to each $[D] \in \mathcal{D}(\vec{B})$ a linear operator*

$$F([D]) : V^{\otimes |\partial_{\text{in}} D|} \rightarrow V^{\otimes |\partial_{\text{out}} D|}$$

that respects the composition and tensor product (Def.4.12), i.e.

$$F([D_1] \circ [D_2]) = F([D_1]) \circ F([D_2]), \quad F([D_1] \otimes [D_2]) = F([D_1]) \otimes F([D_2]),$$

and whose values at elementary classes $[D] \in \mathcal{D}(\vec{B})$, defined in Def.4.13 using Fig.7, are:

- (RT1) (*identity*) if D is as in Fig.7(I) or (VI), then $F([D]) = \text{id} : V \rightarrow V$;
- (RT2) (*cup*) if D is as in Fig.7(II) or (VII), then $F([D]) : \mathbb{C} \rightarrow V \otimes V$ sends $1 \in \mathbb{C}$ to $\omega \xi_+ \otimes \xi_- - \omega^5 \xi_- \otimes \xi_+$;
- (RT3) (*cap*) if D is as in Fig.7(III) or (VIII), then $F([D]) : V \otimes V \rightarrow \mathbb{C}$ sends $\xi_+ \otimes \xi_-$ to $-\omega^{-5}$ and $\xi_- \otimes \xi_+$ to ω^{-1} while other basis vectors to zero;
- (RT4) (*height exchange*) if D is as in Fig.7(IV), then the map $F([D]) : V \otimes V \rightarrow V \otimes V$ is given on the basis vectors as:

$$(5.7) \quad F([D]) : \xi_i \otimes \xi_j \mapsto \begin{cases} \omega^2 \xi_i \otimes \xi_j, & \text{if } i = j, \\ \omega^{-2}(\xi_+ \otimes \xi_- + (\omega^4 - \omega^{-4})\xi_- \otimes \xi_+), & \text{if } i = + \text{ and } j = -, \\ \omega^{-2} \xi_- \otimes \xi_+, & \text{if } i = - \text{ and } j = +, \end{cases}$$

while if D is as in Fig.7(XI), then the map $F([D]) : V \otimes V \rightarrow V \otimes V$ is given by the inverse of the map in eq.(5.7);

- (RT5) (*positive crossing*) if D is as in Fig.7(V), then $F([D]) : V \otimes V \rightarrow V \otimes V$ is given by the composition (inverse of map in eq.(5.7)) $\circ P$, where $P : V \otimes V \rightarrow V \otimes V$ is the position exchange map

$$(5.8) \quad P(\xi_i \otimes \xi_j) = \xi_j \otimes \xi_i, \quad \forall i, j \in \{1, 2\}.$$

- (RT6) (*negative crossing*) if D is as in Fig.7(X), then $F([D]) : V \otimes V \rightarrow V \otimes V$ is given by the inverse of the map $V \otimes V \rightarrow V \otimes V$ for case (RT5).

Lemma 5.11 (orientation insensitivity; [O02, §3]). *$F([D])$ does not depend on the orientation of components of D .*

To make the situation clear, we emphasize again that the validity of the above proposition and lemma is well-known, whose proofs can be found in the original paper [RT90] or the books [T16] [O02]. One remark is that in these original references the Reshetikhin-Turaev operator invariants are formulated for H-tangles, and the condition (RT4) in the above Prop.5.10 is to adapt such constructions to the setting of VH-tangles.

Our strategy to prove Prop.4.15 is to find a relationship between $G([D])$ and $F([D])$, and then to use Prop.5.10. However, a proof of Prop.4.15 for $G([D])$ is not the only reason we recalled the Reshetikhin-Turaev operator invariant $F([D])$. In fact, we observe that the matrix elements of $F([D])$ are precisely the Bonahon-Wong biangle quantum trace. Note that this fact was not mentioned in the original work [BW11]; although it was already observed in [CL19], we present a proof here, for completeness.

Proposition 5.12 (Bonahon-Wong's biangle quantum trace as matrix elements of F ; [CL19, Thm.5.2]). *Let \vec{B} be a directed biangle, with the underlying biangle being $B = (\Sigma', \mathcal{P}')$, with $\mathfrak{S}' = \Sigma' \setminus \mathcal{P}'$. Let $\omega \in \mathbb{C}^*$. Let $[k, s_0]$ be the VH-isotopy class of a stated VH-tangle in $\mathfrak{S}' \times (-1, 1)$.*

Denote by $[D, s_0] \in \mathcal{D}_s(\vec{B})$ the equivalence class of stated boundary-ordered tangle diagram for $[k, s_0]$ (with slight abuse of notation for s_0). Define the inward and outward basis vectors $\xi_{\text{in}}^{s_0} \in V^{\otimes |\partial_{\text{in}} D|}$ and $\xi_{\text{out}}^{s_0} \in V^{\otimes |\partial_{\text{out}} D|}$ as in Def.4.17, and let $F([D, s_0]) \in \mathbb{C}$ be the matrix element of the linear map $F([D]) : V^{\otimes |\partial_{\text{in}} D|} \rightarrow V^{\otimes |\partial_{\text{out}} D|}$ of Prop.5.10 for these basis vectors, i.e.

$$F([D, s_0]) := \langle \xi_{\text{out}}^{s_0}, F([D]) \xi_{\text{in}}^{s_0} \rangle$$

where $\langle \cdot, \cdot \rangle$ is as defined in Def.4.16. Then $F([D, s_0])$ coincides with the Bonahon-Wong biangle quantum trace $\text{Tr}_B^\omega([k, s_0])$ of Prop.3.19:

$$\text{Tr}_B^\omega([k, s_0]) = F([D, s_0]).$$

Proof. For each $[D] \in \mathcal{D}(\vec{B})$ we define $\widehat{F}([D]) : V^{\otimes |\partial_{\text{in}} D|} \rightarrow V^{\otimes |\partial_{\text{out}} D|}$ as the unique linear map whose matrix element of each state s_0 of D coincides with the number $\text{Tr}_B^\omega([k, s_0])$, where (k, s_0) is a stated VH-tangle in $\mathfrak{S}' \times (-1, 1)$ whose projection to \vec{B} yields a stated boundary-ordered tangle diagram equivalent to (D, s_0) (by a slight abuse of notation for s_0):

$$\langle \xi_{\text{out}}^{s_0}, \widehat{F}([D]) \xi_{\text{in}}^{s_0} \rangle = \text{Tr}_B^\omega([k, s_0]).$$

By Prop.3.5(4), any such (k, s_0) are VH-isotopic to each other. Since the Bonahon-Wong biangle quantum trace is defined on a V-isotopy class of stated tangles, we see that $\widehat{F}([D])$ is well-defined. We will now show that $\widehat{F}([D])$ respects the tensor product and composition operation in $\mathcal{D}(\vec{B})$, and that it has same values with $F([D])$ for elementary classes $[D]$.

First, let $[D_1], [D_2] \in \mathcal{D}(\vec{B})$ and consider the tensor product $[D_1] \otimes [D_2]$ defined as $[D'_1 \cup D'_2]$ with D'_1 and D'_2 as in Def.4.12; we shall show that $\widehat{F}([D_1] \otimes [D_2]) = \widehat{F}([D_1]) \otimes \widehat{F}([D_2])$. Let $k'_1 \cup k'_2$ be a VH-tangle in $\mathfrak{S}' \times (-1, 1)$ whose projection gives a diagram equivalent to $D'_1 \cup D'_2$, where k'_1 corresponds to D'_1 and k'_2 to D'_2 . For any state s_0 of $D'_1 \cup D'_2$, we can write it as $s_0 = s_1 \cup s_2$ for states s_1 and s_2 of D'_1 and D'_2 , so that the inward basis vector (eq.(4.6)) for s_0 is $\xi_{\text{in}}^{s_0} = \xi_{\text{in}}^{s_1} \otimes \xi_{\text{in}}^{s_2} \in V^{\otimes |\partial_{\text{in}} D'_1|} \otimes V^{\otimes |\partial_{\text{in}} D'_2|} \equiv V^{\otimes |\partial_{\text{in}}(D'_1 \cup D'_2)|}$. By construction in Def.4.12, D_1 is vertically higher than D_2 , hence k'_1 has higher elevation than k'_2 . Hence the stated skein $[k'_1 \cup k'_2, s_0] = [k'_1 \cup k'_2, s_1 \cup s_2] \in \mathcal{S}_s^A(B)$ equals the product $[k'_2, s_2][k'_1, s_1]$ of stated skeins. Using the fact that the Bonahon-Wong biangle quantum trace map Tr_B^ω is an algebra homomorphism, we obtain

$$\begin{aligned} \langle \xi_{\text{out}}^{s_0}, \widehat{F}([D_1] \otimes [D_2]) \xi_{\text{in}}^{s_0} \rangle &= \langle \xi_{\text{out}}^{s_0}, \widehat{F}([D'_1 \cup D'_2]) \xi_{\text{in}}^{s_0} \rangle = \text{Tr}_B^\omega([k'_1 \cup k'_2, s_0]) = \text{Tr}_B^\omega([k'_2, s_2][k'_1, s_1]) \\ &= \text{Tr}_B^\omega([k'_2, s_2]) \text{Tr}_B^\omega([k'_1, s_1]) = \langle \xi_{\text{out}}^{s_2}, \widehat{F}([D_2]) \xi_{\text{in}}^{s_2} \rangle \langle \xi_{\text{out}}^{s_1}, \widehat{F}([D_1]) \xi_{\text{in}}^{s_1} \rangle = \langle \xi_{\text{out}}^{s_1}, \widehat{F}([D_1]) \xi_{\text{in}}^{s_1} \rangle \langle \xi_{\text{out}}^{s_2}, \widehat{F}([D_2]) \xi_{\text{in}}^{s_2} \rangle \\ &= \langle \xi_{\text{out}}^{s_1} \otimes \xi_{\text{out}}^{s_2}, \widehat{F}([D_1]) \xi_{\text{in}}^{s_1} \otimes \widehat{F}([D_2]) \xi_{\text{in}}^{s_2} \rangle = \langle \xi_{\text{out}}^{s_0}, (\widehat{F}([D_1]) \otimes \widehat{F}([D_2])) \xi_{\text{in}}^{s_0} \rangle \end{aligned}$$

hence $\widehat{F}([D_1] \otimes [D_2]) = \widehat{F}([D_1]) \otimes \widehat{F}([D_2])$ as desired.

Now let $[D_1], [D_2] \in \mathcal{D}(\vec{B})$, and suppose that $[D_1]$ is composable with $[D_2]$ as in Def.4.12; we shall show $\widehat{F}([D_1] \circ [D_2]) = \widehat{F}([D_1]) \circ \widehat{F}([D_2])$. Denote $[D] = [D_1] \circ [D_2]$ (as in Def.4.12), and let k be a VH-tangle in $\mathfrak{S}' \times (-1, 1)$ whose projection gives a diagram equivalent to D , so that cutting k (Def.3.12) along an internal arc of \vec{B} yields VH-tangles k_1 and k_2 in biangles \vec{B}_1 and \vec{B}_2 giving diagrams equivalent to D_1 and D_2 respectively. Then we have a natural identifications $\partial_{\text{out}} k = \partial_{\text{out}} k_1$, $\partial_{\text{in}} k_1 = \partial_{\text{out}} k_2$ and $\partial_{\text{in}} k = \partial_{\text{in}} k_2$. For each state s_0 of k , by the cutting property (Prop.3.19(1)) we have

$$\text{Tr}_B^\omega([k, s_0]) = \sum_{\text{compatible } s_1, s_2} \text{Tr}_{B_1}^\omega([k_1, s_1]) \text{Tr}_{B_2}^\omega([k_2, s_2]),$$

where the sum is over all states s_1 and s_2 of k_1 and k_2 such that $s_1|_{\partial_{\text{out}} k_1} = s_0|_{\partial_{\text{out}} k}$, $s_1|_{\partial_{\text{in}} k_1} = s_2|_{\partial_{\text{out}} k_2}$ and $s_2|_{\partial_{\text{in}} k_2} = s_0|_{\partial_{\text{in}} k}$. Note

$$\begin{aligned} \langle \xi_{\text{out}}^{s_0}, \widehat{F}([D_1] \circ [D_2]) \xi_{\text{in}}^{s_0} \rangle &= \langle \xi_{\text{out}}^{s_0}, \widehat{F}([D]) \xi_{\text{in}}^{s_0} \rangle = \text{Tr}_B^\omega([k, s_0]) \\ &= \sum_{\text{compatible } s_1, s_2} \text{Tr}_{B_1}^\omega([k_1, s_1]) \text{Tr}_{B_2}^\omega([k_2, s_2]) = \sum_{\text{compatible } s_1, s_2} \langle \xi_{\text{out}}^{s_1}, \widehat{F}([D_1]) \xi_{\text{in}}^{s_1} \rangle \langle \xi_{\text{out}}^{s_2}, \widehat{F}([D_2]) \xi_{\text{in}}^{s_2} \rangle. \end{aligned}$$

On the other hand, note

$$\begin{aligned} \langle \xi_{\text{out}}^{s_0}, (\widehat{F}([D_1]) \circ \widehat{F}([D_2])) \xi_{\text{in}}^{s_0} \rangle &= \langle \xi_{\text{out}}^{s_0}, \widehat{F}([D_1]) (\widehat{F}([D_2]) \xi_{\text{in}}^{s_0}) \rangle \\ &= \langle \xi_{\text{out}}^{s_0}, \widehat{F}([D_1]) (\sum_{\vec{\varepsilon}} \langle \xi^{\vec{\varepsilon}}, \widehat{F}([D_2]) \xi_{\text{in}}^{s_0} \rangle \xi^{\vec{\varepsilon}}) \rangle = \sum_{\vec{\varepsilon}} \langle \xi^{\vec{\varepsilon}}, \widehat{F}([D_2]) \xi_{\text{in}}^{s_0} \rangle \langle \xi_{\text{out}}^{s_0}, \widehat{F}([D_1]) \xi^{\vec{\varepsilon}} \rangle \end{aligned}$$

where the sum is over all sign sequences $\vec{\varepsilon} \in \{+, -\}^{|\partial_{\text{out}} D_2|}$. For each $\vec{\varepsilon}$, denote by s_2 the state of k_2 s.t. $s_2|_{\partial_{\text{in}} k_2} = s_0|_{\partial_{\text{in}} k}$ and $s_2|_{\partial_{\text{out}} k_2} = \vec{\varepsilon}$ (using the horizontal ordering of $\partial_{\text{out}} k_2$), and denote by s_1 the state of k_1 s.t. $s_1|_{\partial_{\text{in}} k_1} = \vec{\varepsilon}$ (using the horizontal ordering of $\partial_{\text{in}} k_1$) and $s_1|_{\partial_{\text{out}} k_1} = s|_{\partial_{\text{out}} k}$. Then this sum over $\vec{\varepsilon}$ can be easily seen to be equal to the above sum over compatible s_1 and s_2 . So we obtain $\widehat{F}([D_1] \circ [D_2]) = \widehat{F}([D_1]) \circ \widehat{F}([D_2])$ as desired.

It remains to check the values on the elementary classes. In case of Prop.5.10(RT1), it is easily seen to match Prop.3.19(2)(I). For the case of Prop.5.10(RT2), the values of $\widehat{F}([D])$ can be read from eq.(3.7) of Lem.3.20(1) with x_1 and x_2 playing the roles of y_2 and y_1 , which match the values of $F([D])$ by inspection. For the case of Prop.5.10(RT3), the values of $\widehat{F}([D])$ can be read from Prop.3.19(2)(II) with y_1 and y_2 playing the roles of x_2 and x_1 , which match the values of $F([D])$. For the case of Prop.5.10(RT4), the values of $F([D])$ match the values of $\widehat{F}([D])$ as in eq.(3.9) of Lem.3.20(2) by inspection. The remaining cases (RT5) and (RT6) can be similarly compared with the items (3) and (4) of Lem.3.20 respectively. \blacksquare

Lemma 5.13 (charge conservation of F). *If s is a state of a tangle diagram D in \vec{B} that is not charge-preserving (defined as in Lem.4.18), then the matrix element $\langle \xi_{\text{out}}^s, F([D]) \xi_{\text{in}}^s \rangle$ is zero.*

This lemma is well known in the operator invariant theory (e.g. [O02]), and can also be directly proven in the style of the above proof of Prop.5.12; that is, investigate the behavior under the tensor product and composition in $\mathcal{D}(\vec{B})$, and then elementary classes $[D]$.

5.5. Gabella biangle factors as twisting of Bonahon-Wong biangle factors. We continue from the last subsection, to compare Bonahon-Wong's biangle quantum trace and Gabella's biangle quantum holonomy. We will see that they are not exactly equal, but differ by the signed order correction amount defined in Def.5.1; more precisely, we need an operator version of it.

Definition 5.14 (signed order correction operator). *Let b be a boundary arc of a directed biangle \vec{B} , given the boundary-orientation (Def.2.1). Let Z be an ordered finite set in the interior of b . Define the signed order correction operator of the ordered subset Z on the boundary arc b as the linear operator*

$$\mathcal{C}(b; Z) : V^{\otimes |Z|} \rightarrow V^{\otimes |Z|}$$

given as follows. Let $Z = \{x_1, \dots, x_{|Z|}\}$, written in the increasing order with respect to the horizontal ordering (Def.3.4) on Z coming from the orientation on b . For each state $s_0 : Z \rightarrow \{+, -\}$, denote by ξ^{s_0} the corresponding basis vector of $V^{\otimes |Z|}$

$$\xi^{s_0} := \xi_{s_0(x_1)} \otimes \xi_{s_0(x_2)} \otimes \cdots \otimes \xi_{s_0(x_{|Z|})} \in V^{\otimes |Z|}.$$

Then the map $\mathcal{C}(b; Z)$ is defined by its values on these basis vectors given as

$$\mathcal{C}(b; Z) \xi^{s_0} := \omega^{\mathcal{C}(b; Z, s_0)} \xi^{s_0}$$

for all states s_0 of Z , where $\mathcal{C}(b; Z, s_0)$ is as defined in Def.5.1.

So $\mathcal{C}(b; Z)$ is diagonal and invertible, with the inverse map given by $\mathcal{C}(b; Z)^{-1} \xi^{s_0} = \omega^{-\mathcal{C}(b; Z, s_0)} \xi^{s_0}$.

The following lemma is also one of the crucial technical tools of the present paper.

Lemma 5.15 (relationship between the operator invariants $G([D])$ and $F([D])$). *For any $[D] \in \mathcal{D}(\vec{B})$,*

$$F([D]) = \omega^{2\text{wr}(D)} \mathcal{C}(b_{\text{out}}; \partial_{\text{out}} D) \circ G([D]) \circ \mathcal{C}(b_{\text{in}}; \partial_{\text{in}} D),$$

where the orderings on $\partial_{\text{out}} D$ and $\partial_{\text{in}} D$ (eq.(4.2)) are the vertical orderings.

Keep in mind that in Lem.5.15, b_{out} and b_{in} are given the boundary-orientations (Def.2.1), instead of the orientations described in Def.4.9.

Proof. We define

$$\widehat{G}([D]) := \omega^{-2\text{wr}(D)} \cdot \mathcal{C}(b_{\text{out}}; \partial_{\text{out}} D)^{-1} \circ F([D]) \circ \mathcal{C}(b_{\text{in}}; \partial_{\text{in}} D)^{-1}$$

The strategy is to show that $\widehat{G}([D])$ respects the composition and tensor product operations of $\mathcal{D}(\vec{B})$, and has same values with $G([D])$ on the elementary classes of $\mathcal{D}(\vec{B})$. First, let $[D_1], [D_2] \in \mathcal{D}(\vec{B})$ and consider the tensor product $[D_1] \otimes [D_2]$ defined as $[D'_1 \cup D'_2]$ with D'_1 and D'_2 as in Def.4.12; we shall show that $\widehat{G}([D_1] \otimes [D_2]) = \widehat{G}([D_1]) \otimes \widehat{G}([D_2])$. Since D'_1 and D'_2 are disjoint, we have $\text{wr}(D'_1 \cup D'_2) = \text{wr}(D'_1) + \text{wr}(D'_2)$. For any state s_0 of $D'_1 \cup D'_2$, we can write it as $s_0 = s_1 \cup s_2$ for states s_1 and s_2 of D'_1 and D'_2 , so that the inward basis vector (eq.(4.6)) for s_0 is given by the tensor product $\xi_{\text{in}}^{s_0} = \xi_{\text{in}}^{s_1} \otimes \xi_{\text{in}}^{s_2} \in V^{\otimes |\partial_{\text{in}} D'_1|} \otimes V^{\otimes |\partial_{\text{in}} D'_2|} = V^{\otimes |\partial_{\text{in}}(D'_1 \cup D'_2)|}$. Note

$$\mathcal{C}(b_{\text{in}}; \partial_{\text{in}}(D'_1 \cup D'_2))^{-1}(\xi_{\text{in}}^{s_0}) = \omega^{-\mathcal{C}(b_{\text{in}}; (\partial_{\text{in}} D'_1) \cup (\partial_{\text{in}} D'_2), s_{1;\text{in}} \cup s_{2;\text{in}})} \xi_{\text{in}}^{s_0},$$

where $s_{1;\text{in}} = s_1|_{\partial_{\text{in}} D'_1}$ and $s_{2;\text{in}} = s_2|_{\partial_{\text{in}} D'_2}$. Observe

$$\begin{aligned} \mathcal{C}(b_{\text{in}}; (\partial_{\text{in}} D'_1) \cup (\partial_{\text{in}} D'_2), s_{1;\text{in}} \cup s_{2;\text{in}}) &= \mathcal{C}(b_{\text{in}}; \partial_{\text{in}} D'_1, s_{1;\text{in}}) + \mathcal{C}(b_{\text{in}}; \partial_{\text{in}} D'_2, s_{2;\text{in}}) \\ &+ \underbrace{\sum_{x \in \partial_{\text{in}} D'_2, y \in \partial_{\text{in}} D'_1} \frac{\text{sgn}(b_{\text{in}}; \vec{x}\vec{y})}{=-1} s_0(x) s_0(y)}_{=-1} \end{aligned}$$

where we used the fact that $\partial_{\text{in}} D'_2$ is vertically and horizontally lower than $\partial_{\text{in}} D'_1$, and that the boundary-orientation on b_{in} is going toward the horizontally decreasing direction. The underbraced part equals $-(\sum_{x \in \partial_{\text{in}} D'_2} s_0(x))(\sum_{y \in \partial_{\text{in}} D'_1} s_0(y))$. Likewise,

$$\mathcal{C}(b_{\text{out}}; \partial_{\text{out}}(D'_1 \cup D'_2))^{-1}(\xi_{\text{in}}^{s_0}) = \omega^{-\mathcal{C}(b_{\text{out}}; (\partial_{\text{out}} D'_1) \cup (\partial_{\text{out}} D'_2), s_{1;\text{out}} \cup s_{2;\text{out}})} \xi_{\text{in}}^{s_0},$$

where $s_{1;\text{out}} = s_1|_{\partial_{\text{out}} D'_1}$ and $s_{2;\text{out}} = s_2|_{\partial_{\text{out}} D'_2}$. Observe

$$\begin{aligned} \mathcal{C}(b_{\text{out}}; (\partial_{\text{out}} D'_1) \cup (\partial_{\text{out}} D'_2), s_{1;\text{out}} \cup s_{2;\text{out}}) &= \mathcal{C}(b_{\text{out}}; \partial_{\text{out}} D'_1, s_{1;\text{out}}) + \mathcal{C}(b_{\text{out}}; \partial_{\text{out}} D'_2, s_{2;\text{out}}) \\ &+ \underbrace{\sum_{x \in \partial_{\text{out}} D'_2, y \in \partial_{\text{out}} D'_1} \frac{\text{sgn}(b_{\text{out}}; \vec{x}\vec{y})}{=1} s_0(x) s_0(y)}_{=1} \end{aligned}$$

where we used the fact that $\partial_{\text{out}} D'_2$ is vertically and horizontally lower than $\partial_{\text{out}} D'_1$, and that the boundary-orientation on b_{out} is going toward the horizontally increasing direction. The underbraced part equals $(\sum_{x \in \partial_{\text{out}} D'_2} s_0(x))(\sum_{y \in \partial_{\text{out}} D'_1} s_0(y))$. Combining these, together with the fact

$$F([D_1] \otimes [D_2])(\xi_{\text{in}}^{s_0}) = F([D_1])(\xi_{\text{in}}^{s_1}) \otimes F([D_2])(\xi_{\text{in}}^{s_2}) = (F([D_1]) \otimes F([D_2]))(\xi_{\text{in}}^{s_0}),$$

one obtains

$$\widehat{G}([D_1] \otimes [D_2])(\xi_{\text{in}}^{s_0}) = \omega^{-(*)_{s_0}} (\widehat{G}([D_1])(\xi_{\text{in}}^{s_1}) \otimes \widehat{G}([D_2])(\xi_{\text{in}}^{s_2})) = \omega^{-(*)_{s_0}} (\widehat{G}([D_1]) \otimes \widehat{G}([D_2]))(\xi_{\text{in}}^{s_0}),$$

where $(*)_{s_0}$ is the sum of the underbraced terms:

$$(*)_{s_0} = -(\sum_{x \in \partial_{\text{in}} D'_2} s_0(x))(\sum_{y \in \partial_{\text{in}} D'_1} s_0(y)) + (\sum_{x \in \partial_{\text{out}} D'_2} s_0(x))(\sum_{y \in \partial_{\text{out}} D'_1} s_0(y)).$$

Notice that s is charge-preserving (as defined in Lem.4.18) iff both s_1 and s_2 are charge-preserving. In this case, we have $(*)_{s_0} = 0$, and hence the sought-for equality $\widehat{G}([D_1] \otimes [D_2])(\xi_{\text{in}}^{s_0}) = (\widehat{G}([D_1]) \otimes \widehat{G}([D_2]))(\xi_{\text{in}}^{s_0})$. In case s_0 is not charge-preserving, then by Lem.5.13 we have $\langle \xi_{\text{out}}^{s_0}, \widehat{G}([D_1] \otimes [D_2])(\xi_{\text{in}}^{s_0}) \rangle = 0$, and since s_1 or s_2 is not charge-preserving, by Lem.5.13 we also have $\langle \xi_{\text{out}}^{s_1}, \widehat{G}([D_1])(\xi_{\text{in}}^{s_1}) \rangle = 0$ or $\langle \xi_{\text{out}}^{s_2}, \widehat{G}([D_2])(\xi_{\text{in}}^{s_2}) \rangle = 0$; therefore, $\langle \xi_{\text{out}}^{s_0}, \widehat{G}([D_1] \otimes [D_2])(\xi_{\text{in}}^{s_0}) \rangle = \langle \xi_{\text{out}}^{s_0}, (\widehat{G}([D_1]) \otimes \widehat{G}([D_2]))(\xi_{\text{in}}^{s_0}) \rangle$ holds, and therefore $\widehat{G}([D_1] \otimes [D_2])(\xi_{\text{in}}^{s_0}) = (\widehat{G}([D_1]) \otimes \widehat{G}([D_2]))(\xi_{\text{in}}^{s_0})$.

Now let $[D_1], [D_2] \in \mathcal{D}(\vec{B})$, and suppose that $[D_1]$ is composable with $[D_2]$ as in Def.4.12; we shall show $\widehat{G}([D_1] \circ [D_2]) = \widehat{G}([D_1]) \circ \widehat{G}([D_2])$. If we denote $[D] = [D_1] \circ [D_2]$ (as in Def.4.12), then by construction we see that $\text{wr}(D) = \text{wr}(D_1) + \text{wr}(D_2)$. Note that there is a bijection between $\partial_{\text{in}} D_1$ and $\partial_{\text{out}} D_2$ that preserves both the horizontal and vertical orderings. Thus we have $\mathcal{C}(b_{\text{in}}; \partial_{\text{in}} D_1) = \mathcal{C}(b_{\text{out}}; \partial_{\text{out}} D_2)^{-1}$, in view of the boundary-orientations on b_{in} and b_{out} . Now, note that $\partial_{\text{in}} D$ is naturally identified with $\partial_{\text{in}} D_2$, while $\partial_{\text{out}} D$ with $\partial_{\text{out}} D_1$. Combining all these, together with the fact $F([D_1] \circ [D_2]) = F([D_1]) \circ F([D_2])$, one obtains the sought-for $\widehat{G}([D_1] \circ [D_2]) = \widehat{G}([D_1]) \circ \widehat{G}([D_2])$.

It remains to check the values on the elementary classes. In case of Prop.5.10(RT1), it is easy to see that both $\widehat{G}([D])$ and $G([D])$ are identity maps (see Prop.4.15(GB1)). For the case of Prop.5.10(RT2), we have $\text{wr}(D) = 0$, $\mathcal{C}(b_{\text{in}}; \partial_{\text{in}} D) = \text{id}$ and $\mathcal{C}(b_{\text{out}}; \partial_{\text{out}} D)(\xi_{\pm} \otimes \xi_{\mp}) = 1$. So, we see that $\widehat{G}([D])(1) = \omega^{-1}(\omega \xi_{+} \otimes \xi_{-} - \omega^5 \xi_{-} \otimes \xi_{+})$ which equals $G([D])$, as desired. For the case of Prop.5.10(RT3), we have

$\text{wr}(D) = 0$, $\mathcal{C}(b_{\text{out}}; \partial_{\text{out}}D) = \text{id}$ and $\mathcal{C}(b_{\text{in}}; \partial_{\text{in}}D)(\xi_{\pm} \otimes \xi_{\mp}) = -1$, so $\widehat{G}([D])$ sends $\xi_+ \otimes \xi_-$ to $\omega(-\omega^{-5})$ and $\xi_- \otimes \xi_+$ to $\omega\omega^{-1}$ while other basis vectors to zero, just as $G([D])$, as desired.

For the case of Prop.5.10(RT4) as in Fig.7(IV), we have $\text{wr}(D) = 0$, and for each state s_0 of D we have $\mathcal{C}(b_{\text{in}}; \partial_{\text{in}}D)(\xi_{\text{in}}^{s_0}) = s_0(x_1)s_0(x_2)$ and $\mathcal{C}(b_{\text{out}}; \partial_{\text{out}}D)(\xi_{\text{out}}^{s_0}) = s_0(y_1)s_0(y_2)$, so from Prop.5.10(RT4) we get

$$\widehat{G}([D]) : \xi_i \otimes \xi_j \mapsto \begin{cases} \omega^{-1}\omega^{-1}(\omega^2 \xi_i \otimes \xi_j), & \text{if } i = j, \\ \omega\omega(\omega^{-2}(\xi_+ \otimes \xi_- + (\omega^4 - \omega^{-4})\xi_- \otimes \xi_+)), & \text{if } i = + \text{ and } j = -, \\ \omega\omega(\omega^{-2}\xi_- \otimes \xi_+), & \text{if } i = - \text{ and } j = +. \end{cases}$$

which equals $G([D])$ as in Prop.4.15(GB4). For the case as in Fig.7(XI), we have $\text{wr}(D) = 0$, and for each state s_0 of D we have we have $\mathcal{C}(b_{\text{in}}; \partial_{\text{in}}D)(\xi_{\text{in}}^{s_0}) = -s_0(x_1)s_0(x_2)$ and $\mathcal{C}(b_{\text{out}}; \partial_{\text{out}}D)(\xi_{\text{out}}^{s_0}) = -s_0(y_1)s_0(y_2)$, so one can observe that $\widehat{G}([D])$ coincides with the inverse of the map $\widehat{G}([D])$ for the case of Fig.7(IV), hence also matches $G([D])$ as in Prop.4.15(GB4). For the case of Prop.5.10(RT5), we have $\text{wr}(D) = 1$. For each state s_0 of D , note $\mathcal{C}(b_{\text{in}}; \partial_{\text{in}}D)(\xi_{\text{in}}^{s_0}) = s_0(x_1)s_0(x_2)$ and $\mathcal{C}(b_{\text{out}}; \partial_{\text{out}}D)(\xi_{\text{out}}^{s_0}) = -s_0(y_1)s_0(y_2)$. So, from Prop.5.10(RT5) we get

$$\widehat{G}([D]) : \xi_i \otimes \xi_j \mapsto \begin{cases} \omega^{-2}\omega\omega^{-1}(\omega^{-2}\xi_i \otimes \xi_j), & \text{if } i = j, \\ \omega^{-2}\omega^{-1}\omega(\omega^2 \xi_- \otimes \xi_+), & \text{if } i = + \text{ and } j = -, \\ \omega^{-2}\omega^{-1}\omega(\omega^2(\xi_+ \otimes \xi_- + (\omega^{-4} - \omega^4)\xi_- \otimes \xi_+)), & \text{if } i = - \text{ and } j = +. \end{cases}$$

which equals $G([D])$ as in Prop.4.15(GB5), as desired. For the case of Prop.5.10(RT6), we have $\text{wr}(D) = -1$. For each state s_0 of D , note $\mathcal{C}(b_{\text{in}}; \partial_{\text{in}}D)(\xi_{\text{in}}^{s_0}) = s_0(x_1)s_0(x_2)$ and $\mathcal{C}(b_{\text{out}}; \partial_{\text{out}}D)(\xi_{\text{out}}^{s_0}) = -s_0(y_1)s_0(y_2)$. So one can observe that the map $\widehat{G}([D])$ coincides with the inverse of the map $\widehat{G}([D])$ for the case of Prop.5.10(RT5), and hence equals $G([D])$ as in Prop.4.15(GB6), as desired. \blacksquare

The above proof of Lem.5.15 in particular provides a proof of Prop.4.15, the well-definedness of the invariant $G([D])$, as promised. Now, putting together Def.4.17, Lem.5.15 and Prop.5.12, we obtain a precise relationship between the biangle factors of the two sides.

Lemma 5.16 (relationship between Gabella's biangle quantum holonomy and Bonahon-Wong's biangle quantum trace). *Let \widehat{B} , $B = (\Sigma', \mathcal{P}')$, $\mathfrak{S}' = \Sigma' \setminus \mathcal{P}'$ and ω be as in Prop.5.12. If $[D, s_0] \in \mathcal{D}_s(\widehat{B})$ is the equivalence class of stated boundary-ordered tangle diagram for the VH-isotopy class $[k, s_0]$ of stated VH-tangles in $\mathfrak{S}' \times (-1, 1)$, one has*

$$(5.9) \quad \text{Tr}_B^\omega([k, s_0]) = \omega^{2\text{wr}(D)} \omega^{\partial\mathcal{C}_B(k, s_0)} \text{TrHol}_B^\omega([D, s_0])$$

where $\partial\mathcal{C}_B(k, s_0)$ is the boundary signed order correction amount defined in Def.5.1.

Proof. Note

$$\begin{aligned} & \text{Tr}_B^\omega([k, s_0]) \stackrel{\text{Prop.5.12}}{=} \langle \xi_{\text{out}}^s, F([D])\xi_{\text{in}}^s \rangle \stackrel{\text{Lem.5.15}}{=} \langle \xi_{\text{out}}^s, \omega^{2\text{wr}(D)} \mathcal{C}(b_{\text{out}}; \partial_{\text{out}}D) \circ G([D]) \circ \mathcal{C}(b_{\text{in}}; \partial_{\text{in}}D)\xi_{\text{in}}^s \rangle \\ & \stackrel{\text{Def.5.14}}{=} \langle \xi_{\text{out}}^s, \omega^{2\text{wr}(D)} \mathcal{C}(b_{\text{out}}; \partial_{\text{out}}D) \circ G([D])(\omega^{(*)_1}\xi_{\text{in}}^s) \rangle \quad (\text{with } (*)_1 = \mathcal{C}(b_{\text{in}}; \partial_{\text{in}}D, s_{\text{in}}), \text{ where } s_{\text{in}} = s|_{\partial_{\text{in}}D}) \\ & = \omega^{2\text{wr}(D)+(*)_1} \langle \xi_{\text{out}}^s, \mathcal{C}(b_{\text{out}}; \partial_{\text{out}}D)(\sum_{\vec{\varepsilon}} \langle \xi^{\vec{\varepsilon}}, G([D])\xi_{\text{in}}^s \rangle \xi^{\vec{\varepsilon}}) \rangle \\ & \stackrel{\text{Def.5.14}}{=} \omega^{2\text{wr}(D)+(*)_1} \langle \xi_{\text{out}}^s, (\sum_{\vec{\varepsilon}} \langle \xi^{\vec{\varepsilon}}, G([D])\xi_{\text{in}}^s \rangle \omega^{\mathcal{C}(b_{\text{out}}; \partial_{\text{out}}D, \vec{\varepsilon})} \xi^{\vec{\varepsilon}}) \rangle \quad (\text{with respect to horizontal ordering on } \partial_{\text{out}}D) \\ & \stackrel{\text{Def.4.16}}{=} \omega^{2\text{wr}(D)+\mathcal{C}(b_{\text{in}}; \partial_{\text{in}}D, s_{\text{in}})} \langle \xi_{\text{out}}^s, G([D])\xi_{\text{in}}^s \rangle \omega^{\mathcal{C}(b_{\text{out}}; \partial_{\text{out}}D, s_{\text{out}})} \quad (\text{where } s_{\text{out}} = s|_{\partial_{\text{out}}D}) \\ & \stackrel{\text{Def.4.17}}{=} \omega^{2\text{wr}(D)} \omega^{\mathcal{C}(b_{\text{in}}; \partial_{\text{in}}D, s_{\text{in}})+\mathcal{C}(b_{\text{out}}; \partial_{\text{out}}D, s_{\text{out}})} \text{TrHol}_B^\omega([D, s]). \quad \blacksquare \end{aligned}$$

One consequence of the above Lem.5.16 is a proof of Lem.4.18. Before proving Lem.4.18(1), the direction independence of Gabella biangle quantum holonomy, the TrHol_B^ω in the right hand side of eq.(5.9) should really be written as $\text{TrHol}_{\widehat{B}}^\omega$. However, since all other factors appearing in eq.(5.9) are independent of choice of direction on B , it follows that so is $\text{TrHol}_{\widehat{B}}^\omega([D, s_0])$, as asserted in Lem.4.18(1). For Lem.4.18(2), the charge conservation property of Gabella biangle quantum holonomy $\text{TrHol}_{\widehat{B}}^\omega$, first recall Lem.5.13 which is the charge conservation property of the Bonahon-Wong biangle quantum trace Tr_B^ω . Note that the remaining factors in eq.(5.9) are powers of ω , hence nonzero. Hence the sought-for charge conservation property of $\text{TrHol}_{\widehat{B}}^\omega$ follows.

5.6. Finishing the proof of main theorem. We are only left to assemble everything established so far, to prove the sought-for term-by-term equality eq.(5.2). We first exclude the easy case. For each edge e of $\widehat{\Delta}$, let $b_e(J)$ to be the net sum of signs assigned by J to the junctures lying over e . If there is a biangle B_i (whose sides are e_i and e'_i) such that $b_{e_i}(J) \neq b_{e'_i}(J)$, then by the charge conservation properties (Lem.4.18(2) and Lem.5.13) it follows that the biangle factors $\text{Tr}_{B_i}^\omega([L_i, J|_{\partial L_i}])$ and $\text{TrHol}_{B_i}^\omega([L_i, J|_{\partial L_i}])$ are zero, hence both sides of eq.(5.2) are zero. From now on, we may assume

$$(5.10) \quad b_{e_i}(J) = b_{e'_i}(J) \quad \text{for all } i = 1, \dots, n.$$

The triangle part of the left hand side $\text{BW}_{\widehat{\Delta}}^\omega(K'; J) = \text{BW}_{\widehat{\Delta}}^\omega(K; J)$ is

$$\begin{aligned} \bigotimes_{j=1}^m \text{Tr}_{\widehat{t}_j}^\omega([K_j, J|_{\partial K_j}]) &= \omega^{\sum_{j=1}^m \text{dev}_{\widehat{t}_j}(K_j, J|_{\partial K_j})} \prod_{j=1}^m \left[\text{Tr}_{\widehat{t}_j}^\omega([k_{j,1}, J|_{\partial k_{j,1}}]) \cdots \text{Tr}_{\widehat{t}_j}^\omega([k_{j,l_j}, J|_{\partial k_{j,l_j}}]) \right]_{\text{Weyl}} && (\because \text{eq.(3.10)} \\ & && \text{and Def.5.5}) \\ &= \omega^{\sum_{j=1}^m \text{dev}_{\widehat{t}_j}(K_j, J|_{\partial K_j})} \left[\prod_{j=1}^m \text{Tr}_{\widehat{t}_j}^\omega([k_{j,1}, J|_{\partial k_{j,1}}]) \cdots \text{Tr}_{\widehat{t}_j}^\omega([k_{j,l_j}, J|_{\partial k_{j,l_j}}]) \right]_{\text{Weyl}} \\ &= \omega^{\sum_{j=1}^m \text{dev}_{\widehat{t}_j}(K_j, J|_{\partial K_j})} \left[\prod_{i=1}^n \widehat{Z}_{e_i}^{b_{e_i}(J)} \right]_{\text{Weyl}} && (\because \text{Prop.3.15(2)(a), Def.4.23(G1),} \\ & && \text{eq.(5.10), Def.3.8}) \\ &= \omega^{\sum_{j=1}^m \text{dev}_{\widehat{t}_j}(K_j, J|_{\partial K_j})} \widehat{Z}_{\widetilde{K}^J} && (\because \text{Def.4.23(G1)}) \\ &= \omega^{-4 \sum_{j=1}^m \text{wr}_{\widehat{t}_j}(\widetilde{K}_j^J)} \omega^{2 \sum_{j=1}^m \text{wr}(K_j)} \omega^{\sum_{j=1}^m \partial \mathcal{C}_{\widehat{t}_j}(K_j, J|_{\partial K_j})} \widehat{Z}_{\widetilde{K}^J} && (\because \text{Lem.5.9}) \end{aligned}$$

Note $\omega^{-4 \sum_{j=1}^m \text{wr}_{\widehat{t}_j}(\widetilde{K}_j^J)} = \omega^{-4 \text{wr}_{\widehat{\Delta}}(\widetilde{K}^J)} = q^{-\text{wr}_{\widehat{\Delta}}(\widetilde{K}^J)}$, in view of Def.4.23(G2). From Lem.5.16, we see that the biangle part of $\text{BW}_{\widehat{\Delta}}^\omega(K; J)$ is as follows, if we let D_i be the tangle diagram in B_i for the tangle L_i in the thickening of B_i :

$$\prod_{i=1}^n \text{Tr}_{B_i}^\omega([L_i, J|_{\partial L_i}]) = \omega^{2 \sum_{i=1}^n \text{wr}(D_i)} \omega^{\sum_{i=1}^n \partial \mathcal{C}_{B_i}(L_i, J|_{\partial L_i})} \prod_{i=1}^n \text{TrHol}_{B_i}^\omega([L_i, J|_{\partial L_i}]).$$

Now observe $\sum_{j=1}^m \text{wr}(K_j) + \sum_{i=1}^n \text{wr}(D_i) = \text{wr}(K)$, and also:

Lemma 5.17. $\sum_{j=1}^m \partial \mathcal{C}_{\widehat{t}_j}(K_j, J|_{\partial K_j}) + \sum_{i=1}^n \partial \mathcal{C}_B(L_i, J|_{\partial L_i}) = \partial \mathcal{C}_{(\Sigma, \mathcal{P})}(K, s)$.

Proof. Each edge of $\widehat{\Delta}$ that is not a boundary arc of (Σ, \mathcal{P}) appear as a side of exactly one triangle \widehat{t}_j and one biangle B_i , with opposite orientations, hence the corresponding terms cancel each other. What remain are the boundary signed order correction amounts for the boundary arcs of (Σ, \mathcal{P}) , which form exactly $\partial \mathcal{C}_{(\Sigma, \mathcal{P})}(K, s)$. ■

Combining everything,

$$\begin{aligned} \text{BW}_{\widehat{\Delta}}^\omega(K; J) &= \left(\prod_{i=1}^n \text{Tr}_{B_i}^\omega([L_i, J|_{\partial L_i}]) \right) \left(\bigotimes_{j=1}^m \text{Tr}_{\widehat{t}_j}^\omega([K_j, J|_{\partial K_j}]) \right) && (\text{eq.(3.11)}) \\ &= \omega^{2 \text{wr}(K)} \omega^{\partial \mathcal{C}_{(\Sigma, \mathcal{P})}(K, s)} \left(\prod_{i=1}^n \text{TrHol}_{B_i}^\omega([L_i, J|_{\partial L_i}]) \right) q^{-\text{wr}_{\widehat{\Delta}}(\widetilde{K}^J)} \widehat{Z}_{\widetilde{K}^J} \\ &= \omega^{2 \text{wr}(K)} \omega^{\partial \mathcal{C}_{(\Sigma, \mathcal{P})}(K, s)} \underline{\underline{\Omega}}(\widetilde{K}^J; \omega) \widehat{Z}_{\widetilde{K}^J} && (\text{eq.(4.7)}) \end{aligned}$$

proving the term-by-term equality eq.(5.2), hence the main theorem of the paper.

REFERENCES

- [A16] D.G.L. Allegretti, *The Geometry of Cluster Varieties from Surfaces*, Ph.D. thesis, Yale University, 2016. arXiv: 1606.07788
- [AK17] D.G.L. Allegretti and H. Kim, *A duality map for quantum cluster varieties from surfaces*, Adv. Math. **306** (2017), 1164–1208. arXiv:1509.01567. MR3581328
- [BW11] F. Bonahon and H. Wong, *Quantum traces for representations of surface groups in SL₂(C)*, Geom. Topol. **15** (2011), 1569–1615. MR2851072
- [BW17] F. Bonahon and H. Wong, *Representations of the Kauffman bracket skein algebra II: Punctured surfaces*, Algebr. Geom. Topol. **17** (2017), 3399–3434.
- [BG02] K.A. Brown and K.R. Goodearl, *Lectures on Algebraic Quantum Groups*, Advanced Courses in Mathematics, CRM Barcelona, Birkhäuser Verlag, Basel, 2002.
- [CF99] L.O. Chekhov and V.V. Fock, *Quantum Teichmüller spaces*, Theoret. Math. Phys. **120** (1999), 511–528. arXiv:math/9908165
- [CKKO20] S. Cho, H. Kim, H. Kim and D. Oh, *Laurent positivity of quantized canonical bases for quantum cluster varieties from surface*, Commun. Math. Phys. **373**(2) (2020), 655–705. arXiv:1710.06217.
- [C95] P. M. Cohn, *Skew Fields: Theory of General Division Rings*, Encyclopedia of Mathematics and its Applications **57**, Cambridge University Press, Cambridge, 1995.
- [CL19] F. Costantino and T.T.Q. Lê, *Stated skein algebras of surfaces*, arXiv:1907.11400

- [F97] V.V. Fock, *Dual Teichmüller spaces*, arXiv:dg-ga/9702018
- [FG06] V.V. Fock and A.B. Goncharov, *Moduli spaces of local systems and higher Teichmüller theory*, Publ. Math. Inst. Hautes Études Sci. **103** (2006), 1–211. arXiv:math/0311149v4. MR2233852
- [FG07] V.V. Fock and A.B. Goncharov, *Dual Teichmüller and lamination spaces*, in *Handbook of Teichmüller theory. Vol. I*, IRMA Lect. Math. Theor. Phys. **11**, Eur. Math. Soc., Zürich, 2007, pp. 647–684. arXiv:math/0510312
- [FG09] V.V. Fock and A.B. Goncharov, *The quantum dilogarithm and representations of quantum cluster varieties*, Invent. Math. **175** (2009), 223–286. MR2470108
- [FST08] S. Fomin, M. Shapiro, and D. Thurston, *Cluster algebras and triangulated surfaces. Part I: cluster complexes*, Acta Math. **201**(1) (2008), 83–146. arXiv:math/0608367 MR2448067
- [G17] M. Gabella, *Quantum Holonomies from Spectral Networks and Framed BPS States*, Commun. Math. Phys. **351**(2) (2017), 563–598. arXiv:1603.05258 MR3613514
- [GLM15] D. Galakhov, P. Longhi, and G.W. Moore, *Spectral networks with spin*, Commun. Math. Phys. **340**(1) (2015), 171–232. arXiv:1408.0207
- [G12] D. Gaiotto, *Surface Operators in $\mathcal{N} = 2$ 4d Gauge Theories*, J. High Energy Phys. **1211** (2012), 090. arXiv:0911.1316
- [GMN12] D. Gaiotto, G.W. Moore, and A. Neitzke, *Wall-crossing in coupled 2d-4d systems*, J. High Energy Phys. **12** (2012), 082. arXiv:1103.2598
- [GMN13] D. Gaiotto, G.W. Moore, and A. Neitzke, *Spectral networks*, Ann. Henri Poincaré **14**(7) (2013), 1643–1731. arXiv:1204.4824 MR3115984
- [GMN14] D. Gaiotto, G.W. Moore, and A. Neitzke, *Spectral networks and snakes*, Ann. Henri Poincaré **15**(1) (2014), 61–141. arXiv:1209.0866 MR3147409
- [GHK15] M. Gross, P. Hacking, and S. Keel, *Birational geometry of cluster algebras*, Algebraic Geom. **2**(2) (2015), 137–175. arXiv:1309.2573
- [H10] C. Hiatt, *Quantum traces in quantum Teichmüller theory*, Algebr. Geom. Topol. **10**(3) (2010), 1245–1283. MR2661526
- [K98] R.M. Kashaev, *Quantization of Teichmüller spaces and the quantum dilogarithm*, Lett. Math. Phys. **43** (1998), 105–115. arXiv:q-alg/9705021
- [KS] H. Kim and C. Scarinci, *Quantum moduli spaces of $(2+1)$ -gravity*, in preparation
- [KSv1] H. Kim and M. Son, *SL_2 quantum trace in quantum Teichmüller theory via writhe*, version 1, arXiv:1812.11618v1
- [L17] T.T.Q. Lê, *Quantum Teichmüller spaces and quantum trace map*, J. Inst. Math. Jussieu (2017), 1–43. arXiv:1511.06054
- [L18] T.T.Q. Lê, *Triangular decomposition of skein algebras*, Quantum Topol. **9** (2018), 591–632. arXiv:1609.04987
- [L09] X. Liu, *The quantum Teichmüller space as a noncommutative algebraic object*, J. Knot Theor. Ramif. **18**(5) (2009), 705–726. arXiv:math/0408361 MR2527682
- [O02] T. Ohtsuki, *Quantum Invariants : A study of knots, 3-manifolds and their sets*, Series on Knots and Everything **29**, World Sci. Publ., River Edge, NJ, 2002. MR1881401
- [P12] R.C. Penner, *Decorated Teichmüller theory*, Eur. Math. Soc. Publ., Zürich, 2012.
- [P99] J. Przytycki, *Fundamentals of Kauffman bracket skein modules*, Kobe J. Math. **16** (1999) 45–66.
- [PS00] J. Przytycki and A. Sikora, *On the skein algebras and $SL_2(\mathbb{C})$ -character varieties*, Topology **39** (2000), 115–148.
- [RT90] N.Y. Reshetikhin and V.G. Turaev, *Ribbon graphs and their invariants derived from quantum groups*, Comm. Math. Phys. **127**(1) (1990), 1–26.
- [S20a] L. Shen, *Duals of semisimple Poisson-Lie groups and cluster theory of moduli spaces of G-local systems*, arXiv:2003.07901
- [S20b] M. Son, *Quantum coordinate change map for Chekhov-Fock square root algebras*, Master’s thesis, Ewha Womans University, 2020.
- [T80] W.P. Thurston, *The geometry and topology of 3-manifolds*, lecture notes, Princeton University, 1980, available at <http://library.msri.org/books/gt3m>
- [T91] V.G. Turaev, “Algebra of loops on surfaces, algebra of knots, and quantization” in *Braid group, knot theory and statistical mechanics*, Adv. Ser. Math. Phys. **9**, World Sci. Publ., Teaneck, NJ, 1989. pp59–95.
- [T16] V.G. Turaev, *Quantum invariants of knots and 3-manifolds*, Third edition, De Gruyter Studies in Mathematics, **18**, De Gruyter 2016.

Email address: hyunkyukim@ewha.ac.kr, hyunkyu87@gmail.com

Email address: letu@math.gatech.edu

Email address: mirison@iu.edu

DEPARTMENT OF MATHEMATICS, EWHA WOMANS UNIVERSITY, 52 EWHAYEODAE-GIL, SEODAEMUN-GU, SEOUL 03760, REPUBLIC OF KOREA

SCHOOL OF MATHEMATICS, 686 CHERRY STREET, GEORGIA TECH, ATLANTA, GA 30332, USA

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY BLOOMINGTON, RAWLES HALL, 831 EAST 3RD ST., BLOOMINGTON, IN 47405, USA